

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
10 April 2003 (10.04.2003)

PCT

(10) International Publication Number  
WO 03/029459 A2

- (51) International Patent Classification<sup>7</sup>: C12N 15/11, 37083 Göttingen (DE). RAUHUT, Reinhard [DE/DE]; A61K 48/00 Merkelstrasse 16, 37085 Göttingen (DE).
- (21) International Application Number: PCT/EP02/10881 (74) Agents: WEICKMANN, Franz, Albert et al.; Weickmann & Weickmann, Postfach 860 820, 81635 München (DE).
- (22) International Filing Date: 27 September 2002 (27.09.2002) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
01123453.1 28 September 2001 (28.09.2001) EP  
02006712.0 22 March 2002 (22.03.2002) EP  
02016772.2 26 July 2002 (26.07.2002) EP
- (71) Applicant (*for all designated States except US*): MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN E.V. [DE/DE]; Hofgartenstrasse 8, 80539 München (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): TUSCHL, Thomas [DE/DE]; Keplerstrasse 9, 37085 Göttingen (DE). LAGOS-QUINTANA, Mariana [MX/DE]; Kurze Geismarstrasse 13, 37073 Göttingen (DE). LENDECKEL, Winfried [DE/DE]; Blinde Gasse 52, 37318 Hohengandem (DE). MEYER, Jutta [DE/DE]; Sternstrasse 16,
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: MICRORNA MOLECULES

(57) Abstract: In *Caenorhabditis elegans*, *lin-4* and *let-7* encode 22- and 21 -nucleotide RNAs, respectively, that function as key regulators of developmental timing. Because the appearance of these short RNAs is regulated during development, they are also referred to as "small temporal RNAs" (stRNAs). We show that many more 21- and 22-nt expressed RNAs, termed microRNAs, (miRNAs), exist in invertebrates and vertebrates, and that some of these novel RNAs, similar to *let-7* stRNA, are also highly conserved. This suggests that sequence-specific post-transcriptional regulatory mechanisms mediated by small RNAs are more general than previously appreciated.

WO 03/029459 A2

**MicroRNA molecules****Description**

5

The present invention relates to novel small expressed (micro)RNA molecules associated with physiological regulatory mechanisms, particularly in developmental control.

10

In *Caenorhabditis elegans*, *lin-4* and *let-7* encode 22- and 21-nucleotide RNAs, respectively (1, 2), that function as key regulators of developmental timing (3-5). Because the appearance of these short RNAs is regulated during development, they are also referred to as "microRNAs" (miRNAs) or small temporal RNAs (stRNAs) (6). *lin-4* and *let-21* are the only known miRNAs to date.

15

Two distinct pathways exist in animals and plants in which 21- to 23-nucleotide RNAs function as post-transcriptional regulators of gene expression. Small interfering RNAs (siRNAs) act as mediators of sequence-specific mRNA degradation in RNA interference (RNAi) (7-11) whereas miRNAs regulate developmental timing by mediating sequence-specific repression of mRNA translation (3-5). siRNAs and miRNAs are excised from double-stranded RNA (dsRNA) precursors by Dicer (12, 13, 29), a multidomain RNase III protein, thus producing RNA species of similar size. However, siRNAs are believed to be double-stranded (8, 11, 12), while miRNAs are single-stranded (6).

20

We show that many more short, particularly 21- and 22-nt expressed RNAs, termed microRNAs (miRNAs), exist in invertebrates and vertebrates, and that some of these novel RNAs, similar to *let-7* RNA (6), are also highly conserved. This suggests that sequence-specific post-transcriptional

25

30

regulatory mechanisms mediated by small RNAs are more general than previously appreciated.

The present invention relates to an isolated nucleic acid molecule comprising:

5

(a) a nucleotide sequence as shown in Table 1, Table 2, Table 3 or Table 4

10

(b) a nucleotide sequence which is the complement of (a),

(c) a nucleotide sequence which has an identity of at least 80%, preferably of at least 90% and more preferably of at least 99%, to a sequence of (a) or (b) and/or

15

(d) a nucleotide sequence which hybridizes under stringent conditions to a sequence of (a), (b) and/or (c).

20

In a preferred embodiment the invention relates to miRNA molecules and analogs thereof, to miRNA precursor molecules and to DNA molecules encoding miRNA or miRNA precursor molecules.

Preferably the identity of sequence (c) to a sequence of (a) or (b) is at least 90%, more preferably at least 95%. The determination of identity (percent) may be carried out as follows:

25

$$I = n : L$$

30

wherein I is the identity in percent, n is the number of identical nucleotides between a given sequence and a comparative sequence as shown in Table 1, Table 2, Table 3 or Table 4 and L is the length of the comparative sequence. It should be noted that the nucleotides A, C, G and U as depicted in Tables 1, 2, 3 and 4 may denote ribonucleotides,

- 3 -

deoxyribonucleotides and/or other nucleotide analogs, e.g. synthetic non-naturally occurring nucleotide analogs. Further nucleobases may be substituted by corresponding nucleobases capable of forming analogous H-bonds to a complementary nucleic acid sequence, e.g. U may be substituted by T.

Further, the invention encompasses nucleotide sequences which hybridize under stringent conditions with the nucleotide sequence as shown in Table 1, Table 2, Table 3 or Table 4, a complementary sequence thereof or a highly identical sequence. Stringent hybridization conditions comprise washing for 1 h in 1 x SSC and 0.1% SDS at 45°C, preferably at 48°C and more preferably at 50°C, particularly for 1 h in 0.2 x SSC and 0.1% SDS.

The isolated nucleic acid molecules of the invention preferably have a length of from 18 to 100 nucleotides, and more preferably from 18 to 80 nucleotides. It should be noted that mature miRNAs usually have a length of 19-24 nucleotides, particularly 21, 22 or 23 nucleotides. The miRNAs, however, may be also provided as a precursor which usually has a length of 50-90 nucleotides, particularly 60-80 nucleotides. It should be noted that the precursor may be produced by processing of a primary transcript which may have a length of > 100 nucleotides.

The nucleic acid molecules may be present in single-stranded or double-stranded form. The miRNA as such is usually a single-stranded molecule, while the mi-precursor is usually an at least partially self-complementary molecule capable of forming double-stranded portions, e.g. stem- and loop-structures. DNA molecules encoding the miRNA and miRNA precursor molecules. The nucleic acids may be selected from RNA, DNA or nucleic acid analog molecules, such as sugar- or backbone-modified ribonucleotides or deoxyribonucleotides. It should be noted, however, that other nucleic analogs, such as peptide nucleic acids (PNA) or locked nucleic acids (LNA), are also suitable.

- 4 -

In an embodiment of the invention the nucleic acid molecule is an RNA- or DNA molecule, which contains at least one modified nucleotide analog, i.e. a naturally occurring ribonucleotide or deoxyribonucleotide is substituted by a non-naturally occurring nucleotide. The modified nucleotide analog may be located for example at the 5'-end and/or the 3'-end of the nucleic acid molecule.

Preferred nucleotide analogs are selected from sugar- or backbone-modified ribonucleotides. It should be noted, however, that also nucleobase-modified ribonucleotides, i.e. ribonucleotides, containing a non-naturally occurring nucleobase instead of a naturally occurring nucleobase such as uridines or cytidines modified at the 5-position, e.g. 5-(2-amino)propyl uridine, 5-bromo uridine; adenosines and guanosines modified at the 8-position, e.g. 8-bromo guanosine; deaza nucleotides, e.g. 7-deaza-adenosine; O- and N-alkylated nucleotides, e.g. N6-methyl adenosine are suitable. In preferred sugar-modified ribonucleotides the 2'-OH-group is replaced by a group selected from H, OR, R, halo, SH, SR, NH<sub>2</sub>, NHR, NR<sub>2</sub> or CN, wherein R is C<sub>1</sub>-C<sub>6</sub> alkyl, alkenyl or alkynyl and halo is F, Cl, Br or I. In preferred backbone-modified ribonucleotides the phosphoester group connecting to adjacent ribonucleotides is replaced by a modified group, e.g. of phosphothioate group. It should be noted that the above modifications may be combined.

The nucleic acid molecules of the invention may be obtained by chemical synthesis methods or by recombinant methods, e.g. by enzymatic transcription from synthetic DNA-templates or from DNA-plasmids isolated from recombinant organisms. Typically, phage RNA-polymerases are used for transcription, such as T7, T3 or SP6 RNA-polymerases.

The invention also relates to a recombinant expression vector comprising a recombinant nucleic acid operatively linked to an expression control sequence, wherein expression, i.e. transcription and optionally further

- 5 -

processing results in a miRNA-molecule or miRNA precursor molecule as described above. The vector is preferably a DNA-vector, e.g. a viral vector or a plasmid, particularly an expression vector suitable for nucleic acid expression in eukaryotic, more particularly mammalian cells. The recombinant nucleic acid contained in said vector may be a sequence which results in the transcription of the miRNA-molecule as such, a precursor or a primary transcript thereof, which may be further processed to give the miRNA-molecule.

Further, the invention relates to diagnostic or therapeutic applications of the claimed nucleic acid molecules. For example, miRNAs may be detected in biological samples, e.g. in tissue sections, in order to determine and classify certain cell types or tissue types or miRNA-associated pathogenic disorders which are characterized by differential expression of miRNA-molecules or miRNA-molecule patterns. Further, the developmental stage of cells may be classified by determining temporarily expressed miRNA-molecules.

Further, the claimed nucleic acid molecules are suitable for therapeutic applications. For example, the nucleic acid molecules may be used as modulators or targets of developmental processes or disorders associated with developmental dysfunctions, such as cancer. For example, miR-15 and miR-16 probably function as tumor-suppressors and thus expression or delivery of these RNAs or analogs or precursors thereof to tumor cells may provide therapeutic efficacy, particularly against leukemias, such as B-cell chronic lymphocytic leukemia (B-CLL). Further, miR-10 is a possible regulator of the translation of Hox Genes, particularly Hox 3 and Hox 4 (or Scr and Dfd in *Drosophila*).

In general, the claimed nucleic acid molecules may be used as a modulator of the expression of genes which are at least partially complementary to said nucleic acid. Further, miRNA molecules may act as target for

- 6 -

therapeutic screening procedures, e.g. inhibition or activation of miRNA molecules might modulate a cellular differentiation process, e.g. apoptosis.

Furthermore, existing miRNA molecules may be used as starting materials  
5 for the manufacture of sequence-modified miRNA molecules, in order to modify the target-specificity thereof, e.g. an oncogene, a multidrug-resistance gene or another therapeutic target gene. The novel engineered miRNA molecules preferably have an identity of at least 80% to the starting miRNA, e.g. as depicted in Tables 1, 2, 3 and 4. Further, miRNA  
10 molecules can be modified, in order that they are symmetrically processed and then generated as double-stranded siRNAs which are again directed against therapeutically relevant targets.

Furthermore, miRNA molecules may be used for tissue reprogramming  
15 procedures, e.g. a differentiated cell line might be transformed by expression of miRNA molecules into a different cell type or a stem cell.

For diagnostic or therapeutic applications, the claimed RNA molecules are preferably provided as a pharmaceutical composition. This pharmaceutical  
20 composition comprises as an active agent at least one nucleic acid molecule as described above and optionally a pharmaceutically acceptable carrier.

The administration of the pharmaceutical composition may be carried out  
25 by known methods, wherein a nucleic acid is introduced into a desired target cell in vitro or in vivo.

Commonly used gene transfer techniques include calcium phosphate, DEAE-dextran, electroporation and microinjection and viral methods [30,  
30 31, 32, 33, 34]. A recent addition to this arsenal of techniques for the introduction of DNA into cells is the use of cationic liposomes [35].

- 7 -

Commercially available cationic lipid formulations are e.g. Tfx 50 (Promega) or Lipofectamin 2000 (Life Technologies).

5 The composition may be in form of a solution, e.g. an injectable solution, a cream, ointment, tablet, suspension or the like. The composition may be administered in any suitable way, e.g. by injection, by oral, topical, nasal, rectal application etc. The carrier may be any suitable pharmaceutical carrier. Preferably, a carrier is used, which is capable of increasing the efficacy of the RNA molecules to enter the target-cells. Suitable examples  
10 of such carriers are liposomes, particularly cationic liposomes.

Further, the invention relates to a method of identifying novel microRNA-molecules and precursors thereof, in eukaryotes, particularly in vertebrates and more particularly in mammals, such as humans or mice. This method  
15 comprises: ligating 5'- and 3'-adapter-molecules to the end of a size-fractionated RNA-population, reverse transcribing said adapter-ligated RNA-population, and characterizing said reverse transcribed RNA-molecules, e.g. by amplification, concatamerization, cloning and sequencing.

20 A method as described above already has been described in (8), however, for the identification of siRNA molecules. Surprisingly, it was found now that the method is also suitable for identifying the miRNA molecules or precursors thereof as claimed in the present application.

25 Further, it should be noted that as 3'-adaptor for derivatization of the 3'-OH group not only 4-hydroxymethylbenzyl but other types of derivatization groups, such as alkyl, alkyl amino, ethylene glycol or 3'-deoxy groups are suitable.

30 Further, the invention shall be explained in more detail by the following Figures and Examples:



## Figure Legends

Fig. 1A. Expression of *D. melanogaster* miRNAs. Northern blots of total RNA isolated from staged populations of *D. melanogaster* were probed for the indicated miRNAs. The position of 76-nt val-tRNA is also indicated on the blots. 5S rRNA serves as loading control. E, embryo; L, larval stage; P, pupae; A, adult; S2, Schneider-2 cells. It should be pointed out, that S2 cells are polyclonal, derived from an unknown subset of embryonic tissues, and may have also lost some features of their tissue of origin while maintained in culture. miR-3 to miR-6 RNAs were not detectable in S2 cells (data not shown). miR-14 was not detected by Northern blotting and may be very weakly expressed, which is consistent with its cloning frequency. Similar miRNA sequences are difficult to distinguish by Northern blotting because of potential cross-hybridization of probes.

Fig. 1B. Expression of vertebrate miRNAs. Northern blots of total RNA isolated from HeLa cells, mouse kidneys, adult zebrafish, frog ovaries, and S2 cells were probed for the indicated miRNAs. The position of 76-nt val-tRNA is also indicated on the blots. 5S rRNA from the preparations of total RNA from the indicated species is also shown. The gels used for probing of miR-18, miR-19a, miR-30, and miR-31 were not run as far as the other gels (see tRNA marker position). miR-32 and miR-33 were not detected by Northern blotting, which is consistent with their low cloning frequency. Oligodeoxynucleotides used as Northern probes were:

let-7a, 5' TACTATACAACCTACTACCTCAATTTGCC (SEQ ID NO:1);  
 let-7d, 5' ACTATGCAACCTACTACCTCT (SEQ ID NO:2);  
 let-7e, 5' ACTATACAACCTCCTACCTCA (SEQ ID NO:3);  
*D. melanogaster* val-tRNA, 5' TGGTGTTCGCCCCGGGAA (SEQ ID NO:4);  
 miR-1, 5' TGAATGTAAAGAAGTATGGAG (SEQ ID NO:5);  
 miR-2b, 5' GCTCCTCAAAGCTGGCTGTGATA (SEQ ID NO:6);  
 miR-3, 5' TGAGACACACTTTGCCAGTGA (SEQ ID NO:7);  
 miR-4, 5' TCAATGGTTGTCTAGCTTTAT (SEQ ID NO:8);

- 9 -

miR-5, 5' CATATCACAACGATCGTTCCTTT (SEQ ID NO:9);  
miR-6, 5' AAAAAGAACAGCCACTGTGATA (SEQ ID NO:10);  
miR-7, 5' TGGAAGACTAGTGATTTTGTGT (SEQ ID NO:11);  
miR-8, 5' GACATCTTTACCTGACAGTATTA (SEQ ID NO:12);  
5 miR-9, 5' TCATACAGCTAGATAACCAAAGA (SEQ ID NO:13);  
miR-10, 5' ACAAATTCGGATCTACAGGGT (SEQ ID NO:14);  
miR-11, 5' GCAAGAACTCAGACTGTGATG (SEQ ID NO:15);  
miR-12, 5' ACCAGTACCTGATGTAATACTCA (SEQ ID NO:16);  
miR-13a, 5' ACTCGTCAAAATGGCTGTGATA (SEQ ID NO:17);  
10 miR-14, 5' TAGGAGAGAGAAAAGACTGA (SEQ ID NO:18);  
miR-15, 5' TAGCAGCACATAATGGTTTGT (SEQ ID NO:19);  
miR-16, 5' GCCAATATTTACGTGCTGCTA (SEQ ID NO:20);  
miR-17, 5' TACAAGTGCCTTCACTGCAGTA (SEQ ID NO:21);  
miR-18, 5' TATCTGCACTAGATGCACCTTA (SEQ ID NO:22);  
15 miR-19a, 5' TCAGTTTTGCATAGATTTGCACA (SEQ ID NO:23);  
miR-20, 5' TACCTGCACTATAAGCACTTTA (SEQ ID NO:24);  
miR-21, 5' TCAACATCAGTCTGATAAGCTA (SEQ ID NO:25);  
miR-22, 5' ACAGTTCTTCAACTGGCAGCTT (SEQ ID NO:26);  
miR-23, 5' GGAAATCCCTGGCAATGTGAT (SEQ ID NO:27);  
20 miR-24, 5' CTGTTCCTGCTGAACTGAGCCA (SEQ ID NO:28);  
miR-25, 5' TCAGACCGAGACAAGTGCAATG (SEQ ID NO:29);  
miR-26a, 5' AGCCTATCCTGGATTACTTGAA (SEQ ID NO:30);  
miR-27, 5' AGCGGAACTTAGCCACTGTGAA (SEQ ID NO:31);  
miR-28, 5' CTCAATAGACTGTGAGCTCCTT (SEQ ID NO:32);  
25 miR-29, 5' AACCGATTTTCAGATGGTGCTAG (SEQ ID NO:33);  
miR-30, 5' GCTGCAAACATCCGACTGAAAG (SEQ ID NO:34);  
miR-31, 5' CAGCTATGCCAGCATCTTGCCT (SEQ ID NO:35);  
miR-32, 5' GCAACTTAGTAATGTGCAATA (SEQ ID NO:36);  
miR-33, 5' TGCAATGCAACTACAATGCACC (SEQ ID NO:37).

30

Fig. 2. Genomic organization of miRNA gene clusters. The precursor structure is indicated as box and the location of the miRNA within the

- 10 -

precursor is shown in gray; the chromosomal location is also indicated to the right. (A) *D. melanogaster* miRNA gene clusters. (B) Human miRNA gene clusters. The cluster of let-7a-1 and let-7f-1 is separated by 26500 nt from a copy of let-7d on chromosome 9 and 17. A cluster of let-7a-3 and let-7b, separated by 938 nt on chromosome 22, is not illustrated.

Fig. 3. Predicted precursor structures of *D. melanogaster* miRNAs. RNA secondary structure prediction was performed using mfold version 3.1 [28] and manually refined to accommodate G/U wobble base pairs in the helical segments. The miRNA sequence is underlined. The actual size of the stem-loop structure is not known experimentally and may be slightly shorter or longer than represented. Multicopy miRNAs and their corresponding precursor structures are also shown.

Fig. 4. Predicted precursor structures of human miRNAs. For legend, see Fig. 3.

Fig. 5. Expression of novel mouse miRNAs. Northern blot analysis of novel mouse miRNAs. Total RNA from different mouse tissues was blotted and probed with a 5'-radiolabeled oligodeoxynucleotide complementary to the indicated miRNA. Equal loading of total RNA on the gel was verified by ethidium bromide staining prior to transfer; the band representing tRNAs is shown. The fold-back precursors are indicated with capital L. Mouse brains were dissected into midbrain, mb, cortex, cx, cerebellum, cb. The rest of the brain, rb, was also used. Other tissues were heart, ht, lung, lg, liver, lv, colon, co, small intestine, si, pancreas, pc, spleen, sp, kidney, kd, skeletal muscle, sm, stomach, st, H, human Hela SS3 cells. Oligodeoxynucleotides used as Northern probes were:

miR-1a, CTCCATACTTCTTTACATTCCA (SEQ ID NO:38);  
miR-30b, GCTGAGTGTAGGATGTTTACA (SEQ ID NO:39);  
miR-30a-s, GCTTCCAGTCGAGGATGTTTACA (SEQ ID NO:40);  
miR-99b, CGCAAGGTCGGTTCTACGGGTG (SEQ ID NO:41);

- 11 -

miR-101, TCAGTTATCACAGTACTGTA (SEQ ID NO:42);  
 miR-122a, ACAAACACCATTGTCACTCCA (SEQ ID NO:43);  
 miR-124a, TGGCATTACCGCGTGCCTTA (SEQ ID NO:44);  
 miR-125a, CACAGGTTAAAGGGTCTCAGGGA (SEQ ID NO:45);  
 5 miR-125b, TCACAAGTTAGGGTCTCAGGGA (SEQ ID NO:46);  
 miR-127, AGCCAAGCTCAGACGGATCCGA (SEQ ID NO:47);  
 miR-128, AAAAGAGACCGGTTCACTCTGA (SEQ ID NO:48);  
 miR-129, GCAAGCCCAGACCGAAAAAAG (SEQ ID NO:49);  
 miR-130, GCCCTTTTAACATTGCACTC (SEQ ID NO:50);  
 10 miR-131, ACTTTCGGTTATCTAGCTTTA (SEQ ID NO:51);  
 miR-132, ACGACCATGGCTGTAGACTGTTA (SEQ ID NO:52);  
 miR-143, TGAGCTACAGTGCTTCATCTCA (SEQ ID NO:53).

15 Fig.6. Potential orthologs of lin-4 stRNA. (A) Sequence alignment of *C. elegans* lin-4 stRNA with mouse miR-125a and miR-125b and the *D. melanogaster* miR-125. Differences are highlighted by gray boxes. (B) Northern blot of total RNA isolated from staged populations of *D. melanogaster*, probed for miR-125. E, embryo; L, larval stage; P, pupae; A,  
 20 adult; S2, Schneider-2 cells.

Fig. 7. Predicted precursor structures of miRNAs, sequence accession numbers and homology information. RNA secondary structure prediction was performed using mfold version 3.1 and manually refined to  
 25 accommodate G/U wobble base pairs in the helical segments. Dashes were inserted into the secondary structure presentation when asymmetrically bulged nucleotides had to be accommodated. The excised miRNA sequence is underlined. The actual size of the stem-loop structure is not known experimentally and may be slightly shorter or longer than  
 30 represented. Multicopy miRNAs and their corresponding precursor structures are also shown. In cases where no mouse precursors were yet deposited in the database, the human orthologs are indicated. miRNAs

- 12 -

which correspond to *D. melanogaster* or human sequences are included. Published *C. elegans* miRNAs [36, 37] are also included in the table. A recent set of new HeLa cell miRNAs is also indicated [46]. If several ESTs were retrieved for one organism in the database, only those with different precursor sequences are listed. miRNA homologs found in other species are indicated. Chromosomal location and sequence accession numbers, and clusters of miRNA genes are indicated. Sequences from cloned miRNAs were searched against mouse and human in GenBank (including trace data), and against *Fugu rubripes* and *Danio rerio* at [www.jgi.doe.gov](http://www.jgi.doe.gov) and [www.sanger.ac.uk](http://www.sanger.ac.uk), respectively.

EXAMPLE 1: MicroRNAs from *D. melanogaster* and human.

We previously developed a directional cloning procedure to isolate siRNAs after processing of long dsRNAs in *Drosophila melanogaster* embryo lysate (8). Briefly, 5' and 3' adapter molecules were ligated to the ends of a size-fractionated RNA population, followed by reverse transcription, PCR amplification, concatamerization, cloning and sequencing. This method, originally intended to isolate siRNAs, led to the simultaneous identification of 14 novel 20- to 23-nt short RNAs which are encoded in the *D. melanogaster* genome and which are expressed in 0 to 2 h embryos (Table 1). The method was adapted to clone RNAs in a similar size range from HeLa cell total RNA (14), which led to the identification of 19 novel human stRNAs (Table 2), thus providing further evidence for the existence of a large class of small RNAs with potential regulatory roles. According to their small size, we refer to these novel RNAs as microRNAs or miRNAs. The miRNAs are abbreviated as miR-1 to miR-33, and the genes encoding miRNAs are named mir-1 to mir-33. Highly homologous miRNAs are classified by adding a lowercase letter, followed by a dash and a number for designating multiple genomic copies of a mir gene.

- 13 -

The expression and size of the cloned, endogenous short RNAs was also examined by Northern blotting (Fig. 1, Table 1 and 2). Total RNA isolation was performed by acid guanidinium thiocyanate-phenol-chloroform extraction [45]. Northern analysis was performed as described [1], except that the total RNA was resolved on a 15% denaturing polyacrylamide gel, transferred onto Hybond-N+ membrane (Amersham Pharmacia Biotech), and the hybridization and wash steps were performed at 50°C. Oligodeoxynucleotides used as Northern probes were 5'-32P-phosphorylated, complementary to the miRNA sequence and 20 to 25 nt in length.

5S rRNA was detected by ethidium staining of polyacrylamide gels prior to transfer. Blots were stripped by boiling in 0.1% aqueous sodium dodecylsulfate/0.1x SSC (15 mM sodium chloride, 1.5 mM sodium citrate, pH 7.0) for 10 min, and were re-probed up to 4 times until the 21-nt signals became too weak for detection. Finally, blots were probed for val-tRNA as size marker.

For analysis of *D. melanogaster* RNAs, total RNA was prepared from different developmental stages, as well as cultured Schneider-2 (S2) cells, which originally derive from 20-24 h *D. melanogaster* embryos [15] (Fig. 1, Table 1). miR-3 to miR-7 are expressed only during embryogenesis and not at later developmental stages. The temporal expression of miR-1, miR-2 and miR-8 to miR-13 was less restricted. These miRNAs were observed at all developmental stages though significant variations in the expression levels were sometimes observed. Interestingly, miR-1, miR-3 to miR-6, and miR-8 to miR-11 were completely absent from cultured Schneider-2 (S2) cells, which were originally derived from 20-24 h *D. melanogaster* embryos [15], while miR-2, miR-7, miR-12, and miR-13 were present in S2 cells, therefore indicating cell type-specific miRNA expression. miR-1, miR-8, and miR-12 expression patterns are similar to those of lin-4 stRNA in *C. elegans*, as their expression is strongly upregulated in larvae and sustained

- 14 -

to adulthood [16]. miR-9 and miR-11 are present at all stages but are strongly reduced in the adult which may reflect a maternal contribution from germ cells or expression in one sex only.

5 The mir-3 to mir-6 genes are clustered (Fig. 2A), and mir-6 is present as triple repeat with slight variations in the mir-6 precursor sequence but not in the miRNA sequence itself. The expression profiles of miR-3 to miR-6 are highly similar (Table 1), which suggests that a single embryo-specific precursor transcript may give rise to the different miRNAs, or that the  
10 same enhancer regulates miRNA-specific promoters. Several other fly miRNAs are also found in gene clusters (Fig. 2A).

The expression of HeLa cell miR-15 to miR-33 was examined by Northern blotting using HeLa cell total RNA, in addition to total RNA prepared from  
15 mouse kidneys, adult zebrafish, *Xenopus laevis* ovary, and *D. melanogaster* S2 cells (Fig. 1B, Table 2). miR-15 and miR-16 are encoded in a gene cluster (Fig. 2B) and are detected in mouse kidney, fish, and very weakly in frog ovary, which may result from miRNA expression in somatic ovary tissue rather than oocytes. mir-17 to mir-20 are also clustered (Fig. 2B),  
20 and are expressed in HeLa cells and fish, but undetectable in mouse kidney and frog ovary (Fig. 1, Table 2), and therefore represent a likely case of tissue-specific miRNA expression.

The majority of vertebrate and invertebrate miRNAs identified in this study  
25 are not related by sequence, but a few exceptions, similar to the highly conserved let-7 RNA [6], do exist. Sequence analysis of the *D. melanogaster* miRNAs revealed four such examples of sequence conservation between invertebrates and vertebrates. miR-1 homologs are encoded in the genomes of *C. elegans*, *C. briggsae*, and humans, and are  
30 found in cDNAs from zebrafish, mouse, cow and human. The expression of mir-1 was detected by Northern blotting in total RNA from adult zebrafish and *C. elegans*, but not in total RNA from HeLa cells or mouse kidney

- 15 -

(Table 2 and data not shown). Interestingly, while mir-1 and let-7 are expressed both in adult flies (Fig. 1A) [6] and are both undetected in S2 cells, miR-1 is, in contrast to let-7, undetectable in HeLa cells. This represents another case of tissue-specific expression of a miRNA, and indicates that miRNAs may not only play a regulatory role in developmental timing, but also in tissue specification. miR-7 homologs were found by database searches in mouse and human genomic and expressed sequence tag sequences (ESTs). Two mammalian miR-7 variants are predicted by sequence analysis in mouse and human, and were detected by Northern blotting in HeLa cells and fish, but not in mouse kidney (Table 2). Similarly, we identified mouse and human miR-9 and miR-10 homologs by database searches but only detected mir-10 expression in mouse kidney.

The identification of evolutionary related miRNAs, which have already acquired multiple sequence mutations, was not possible by standard bioinformatic searches. Direct comparison of the *D. melanogaster* miRNAs with the human miRNAs identified an 11-nt segment shared between *D. melanogaster* miR-6 and HeLa miR-27, but no further relationships were detected. One may speculate that most miRNAs only act on a single target and therefore allow for rapid evolution by covariation, and that highly conserved miRNAs act on more than one target sequence, and therefore have a reduced probability for evolutionary drift by covariation [6]. An alternative interpretation is that the sets of miRNAs from *D. melanogaster* and humans are fairly incomplete and that many more miRNAs remain to be discovered, which will provide the missing evolutionary links.

lin-4 and let-7 stRNAs were predicted to be excised from longer transcripts that contain approximately 30 base-pair stem-loop structures [1, 6]. Database searches for newly identified miRNAs revealed that all miRNAs are flanked by sequences that have the potential to form stable stem-loop structures (Fig. 3 and 4). In many cases, we were able to detect the predicted, approximately 70-nt precursors by Northern blotting (Fig. 1).



- 16 -

Some miRNA precursor sequences were also identified in mammalian cDNA (EST) databases [27], indicating that primary transcripts longer than 70-nt stem-loop precursors do also exist. We never cloned a 22-nt RNA complementary to any of the newly identified miRNAs, and it is as yet  
5 unknown how the cellular processing machinery distinguishes between the miRNA and its complementary strand. Comparative analysis of the precursor stem-loop structures indicates that the loops adjacent to the base-paired miRNA segment can be located on either side of the miRNA sequence (Fig. 3 and 4), suggesting that the 5' or 3' location of the stem-  
10 closing loop is not the determinant of miRNA excision. It is also unlikely that the structure, length or stability of the precursor stem is the critical determinant as the base-paired structures are frequently imperfect and interspersed by less stable, non-Watson-Crick base pairs such as G/A, U/U, C/U, A/A, and G/U wobbles. Therefore, a sequence-specific recognition  
15 process is a likely determinant for miRNA excision, perhaps mediated by members of the Argonaute (*rde-1/ago1/piwi*) protein family. Two members of this family, *alg-1* and *alg-2*, have recently been shown to be critical for stRNA processing in *C. elegans* [13]. Members of the Argonaute protein family are also involved in RNAi and PTGS. In *D. melanogaster*, these  
20 include *argonaute2*, a component of the siRNA-endonuclease complex (RISC) [17], and its relative *aubergine*, which is important for silencing of repeat genes [18]. In other species, these include *rde-1*, *argonaute1*, and *qde-2*, in *C. elegans* [19], *Arabidopsis thaliana* [20], and *Neurospora crassa* [21], respectively. The Argonaute protein family therefore represents,  
25 besides the RNase III Dicer [12, 13], another evolutionary link between RNAi and miRNA maturation.

Despite advanced genome projects, computer-assisted detection of genes encoding functional RNAs remains problematic [22]. Cloning of expressed,  
30 short functional RNAs, similar to EST approaches (RNomics), is a powerful alternative and probably the most efficient method for identification of such novel gene products [23-26]. The number of functional RNAs has been

- 17 -

widely underestimated and is expected to grow rapidly because of the development of new functional RNA cloning methodologies.

5 The challenge for the future is to define the function and the potential targets of these novel miRNAs by using bioinformatics as well as genetics, and to establish a complete catalogue of time- and tissue-specific distribution of the already identified and yet to be uncovered miRNAs. lin-4 and let-7 stRNAs negatively regulate the expression of proteins encoded by mRNAs whose 3' untranslated regions contain sites of complementarity to  
10 the stRNA [3-5].

Thus, a series of 33 novel genes, coding for 19- to 23-nucleotide microRNAs (miRNAs), has been cloned from fly embryos and human cells. Some of these miRNAs are highly conserved between vertebrates and  
15 invertebrates and are developmentally or tissue-specifically expressed. Two of the characterized human miRNAs may function as tumor suppressors in B-cell chronic lymphocytic leukemia. miRNAs are related to a small class of previously described 21- and 22-nt RNAs (lin-4 and let-7 RNAs), so-called small temporal RNAs (stRNAs), and regulate developmental timing in C.  
20 elegans and other species. Similar to stRNAs, miRNAs are presumed to regulate translation of specific target mRNAs by binding to partially complementary sites, which are present in their 3'-untranslated regions.

Deregulation of miRNA expression may be a cause of human disease, and  
25 detection of expression of miRNAs may become useful as a diagnostic. Regulated expression of miRNAs in cells or tissue devoid of particular miRNAs may be useful for tissue engineering, and delivery or transgenic expression of miRNAs may be useful for therapeutic intervention. miRNAs may also represent valuable drug targets itself. Finally, miRNAs and their  
30 precursor sequences may be engineered to recognize therapeutic valuable targets.

- 18 -

EXAMPLE 2: miRNAs from mouse.

To gain more detailed insights into the distribution and function of miRNAs in mammals, we investigated the tissue-specific distribution of miRNAs in adult mouse. Cloning of miRNAs from specific tissues was preferred over whole organism-based cloning because low-abundance miRNAs that normally go undetected by Northern blot analysis are identified clonally. Also, in situ hybridization techniques for detecting 21-nt RNAs have not yet been developed. Therefore, 19- to 25-nucleotide RNAs were cloned and sequenced from total RNA, which was isolated from 18.5 weeks old BL6 mice. Cloning of miRNAs was performed as follows: 0.2 to .1 mg. of total RNA was separated on a 15% denaturing polyacrylamide gel and RNA of 19- to 25-nt size was recovered. A 5'-phosphorylated 3'-adapter oligonucleotide (5'-pUUUaaccgccaattccagx: uppercase, RNA; lowercase, DNA; p, phosphate; x, 3'-Amino-Modifier C-7, ChemGenes, Ashland, Ma, USA, Cat. No. NSS-1004; SEQ ID NO:54) and a 5'-adapter oligonucleotide (5'-acggaattcctcactAAA: uppercase, RNA; lowercase, DNA; SEQ ID NO:55) were ligated to the short RNAs. RT/PCR was performed with 3'-primer (5'-GACTAGCTGGAATTCGCGGTAAA; SEQ ID NO:56) and 5'-primer (5'-CAGCCAACGGAATTCCTCACTAAA; SEQ ID NO:57). In order to introduce Ban I restriction sites, a second PCR was performed using the primer pair 5'-CAGCCAACAGGCACCGAATTCCTCACTAAA (SEQ ID NO:57) and 5'-GACTAGCTTGGTGCCGAATTCGCGGTAAA (SEQ ID NO:56), followed by concatamerization after Ban I digestion and T4 DNA ligation. Concatamers of 400 to 600 basepairs were cut out from 1.5% agarose gels and recovered by Biotrap (Schleicher & Schuell) electroelution (1x TAE buffer) and by ethanol precipitation. Subsequently, the 3' ends of the concatamers were filled in by incubating for 15 min at 72°C with Taq polymerase in standard PCR reaction mixture. This solution was diluted 3-fold with water and directly used for ligation into pCR2.1 TOPO vectors. Clones were screened for inserts by PCR and 30 to 50 samples were subjected to sequencing. Because RNA was prepared from combining

- 19 -

tissues of several mice, minor sequence variations that were detected multiple times in multiple clones may reflect polymorphisms rather than RT/PCR mutations. Public database searching was used to identify the genomic sequences encoding the approx. 21-nt RNAs. The occurrence of a 20 to 30 basepair fold-back structure involving the immediate upstream or downstream flanking sequences was used to assign miRNAs [36-38].

We examined 9 different mouse tissues and identified 34 novel miRNAs, some of which are highly tissue-specifically expressed (Table 3 and Figure 5). Furthermore, we identified 33 new miRNAs from different mouse tissues and also from human Soas-2 osteosarcoma cells (Table 4). miR-1 was previously shown by Northern analysis to be strongly expressed in adult heart, but not in brain, liver, kidney, lung or colon [37]. Here we show that miR-1 accounts for 45% of all mouse miRNAs found in heart, yet miR-1 was still expressed at a low level in liver and midbrain even though it remained undetectable by Northern analysis. Three copies or polymorphic alleles of miR-1 were found in mice. The conservation of tissue-specific miR-1 expression between mouse and human provides additional evidence for a conserved regulatory role of this miRNA. In liver, variants of miR-122 account for 72% of all cloned miRNAs and miR-122 was undetected in all other tissues analyzed. In spleen, miR-143 appeared to be most abundant, at a frequency of approx. 30%. In colon, miR-142-as, was cloned several times and also appeared at a frequency of 30%. In small intestine, too few miRNA sequences were obtained to permit statistical analysis. This was due to strong RNase activity in this tissue, which caused significant breakdown of abundant non-coding RNAs, e.g. rRNA, so that the fraction of miRNA in the cloned sequences was very low. For the same reason, no miRNA sequences were obtained from pancreas.

30

To gain insights in neural tissue miRNA distribution, we analyzed cortex, cerebellum and midbrain. Similar to heart, liver and small intestine, variants

- 20 -

of a particular miRNA, miR-124, dominated and accounted for 25 to 48% of all brain miRNAs. miR-101, -127, -128, -131, and -132, also cloned from brain tissues, were further analyzed by Northern blotting and shown to be predominantly brain-specific. Northern blot analysis was performed as described in Example 1. tRNAs and 5S rRNA were detected by ethidium staining of polyacrylamide gels prior to transfer to verify equal loading. Blots were stripped by boiling in deionized water for 5 min, and reprobbed up to 4 times until the 21-nt signals became too weak for detection.

miR-125a and miR-125b are very similar to the sequence of *C. elegans* lin-4 stRNA and may represent its orthologs (Fig. 6A). This is of great interest because, unlike let-7 that was readily detected in other species, lin-4 has acquired a few mutations in the central region and thus escaped bioinformatic database searches. Using the mouse sequence miR-125b, we could readily identify its ortholog in the *D. melanogaster* genome. miR-125a and miR-125b differ only by a central diuridine insertion and a U to C change. miR-125b is very similar to lin-4 stRNA with the differences located only in the central region, which is presumed to be bulged out during target mRNA recognition [41]. miR-125a and miR-125b were cloned from brain tissue, but expression was also detected by Northern analysis in other tissues, consistent with the role for lin-4 in regulating neuronal remodeling by controlling lin-14 expression [43]. Unfortunately, orthologs to *C. elegans* lin-14 have not been described and miR-125 targets remain to be identified in *D. melanogaster* or mammals. Finally, miR-125b expression is also developmentally regulated and only detectable in pupae and adult but not in embryo or larvae of *D. melanogaster* (Fig. 6B).

Sequence comparison of mouse miRNAs with previously described miRNA reveals that miR-99b and miR-99a are similar to *D. melanogaster*, mouse and human miR-10 as well as *C. elegans* miR-51 [36], miR-141 is similar to *D. melanogaster* miR-8, miR-29b is similar to *C. elegans* miR-83, and miR-131 and miR-142-s are similar to *D. melanogaster* miR-4 and *C.*

- 21 -

*C. elegans* miR-79 [36]. miR-124a is conserved between invertebrates and vertebrates. In this respect it should be noted that for almost every miRNA cloned from mouse was also encoded in the human genome, and frequently detected in other vertebrates, such as the pufferfish, *Fugu rubripes*, and the zebrafish, *Danio rerio*. Sequence conservation may point to conservation in function of these miRNAs. Comprehensive information about orthologous sequences is listed in Fig. 7.

In two cases both strands of miRNA precursors were cloned (Table 3), which was previously observed once for a *C. elegans* miRNA [36]. It is thought that the most frequently cloned strand of a miRNA precursor represents the functional miRNA, which is miR-30c-s and miR-142-as, s and as indicating the 5' or 3' side of the fold-back structure, respectively.

The mir-142 gene is located on chromosome 17, but was also found at the breakpoint junction of a t(8;17) translocation, which causes an aggressive B-cell leukemia due to strong up-regulation of a translocated MYC gene [44]. The translocated MYC gene, which was also truncated at the first exon, was located only 4-nt downstream of the 3'-end of the miR-142 precursor. This suggests that translocated MYC was under the control of the upstream miR-142 promoter. Alignment of mouse and human miR-142 containing EST sequences indicate an approximately 20 nt conserved sequence element downstream of the mir-142 hairpin. This element was lost in the translocation. It is conceivable that the absence of the conserved downstream sequence element in the putative miR-142/mRNA fusion prevented the recognition of the transcript as a miRNA precursor and therefore may have caused accumulation of fusion transcripts and overexpression of MYC.

miR-155, which was cloned from colon, is excised from the known noncoding BIC RNA [47]. BIC was originally identified as a gene transcriptionally activated by promoter insertion at a common retroviral

- 22 -

integration site in B cell lymphomas induced by avian leukosis virus. Comparison of BIC cDNAs from human, mouse and chicken revealed 78% identity over 138 nucleotides [47]. The identity region covers the miR-155 fold-back precursor and a few conserved boxes downstream of the fold-back sequence. The relatively high level of expression of BIC in lymphoid organs and cells in human, mouse and chicken implies an evolutionary conserved function, but BIC RNA has also been detected at low levels in non-hematopoietic tissues [47].

Another interesting observation was that segments of perfect complementarity to miRNAs are not observed in mRNA sequences or in genomic sequences outside the miRNA inverted repeat. Although this could be fortuitous, based on the link between RNAi and miRNA processing [11, 13, 43] it may be speculated that miRNAs retain the potential to cleave perfectly complementary target RNAs. Because translational control without target degradation could provide more flexibility it may be preferred over mRNA degradation.

In summary, 63 novel miRNAs were identified from mouse and 4 novel miRNAs were identified from human Soas-2 osteosarcoma cells (Table 3 and Table 4), which are conserved in human and often also in other non-mammalian vertebrates. A few of these miRNAs appear to be extremely tissue-specific, suggesting a critical role for some miRNAs in tissue-specification and cell lineage decisions. We may have also identified the fruitfly and mammalian ortholog of *C. elegans* lin-4 stRNA. The establishment of a comprehensive list of miRNA sequences will be instrumental for bioinformatic approaches that make use of completed genomes and the power of phylogenetic comparison in order to identify miRNA-regulated target mRNAs.

**References and Notes**

1. R. C. Lee, R. L. Feinbaum, V. Ambros, *Cell* 75, 843 (1993).
2. B. J. Reinhart et al., *Nature* 403, 901 (2000).
- 5 3. V. Ambros, *Curr. Opin. Genet. Dev.* 10, 428 (2000).
4. E. G. Moss, *Curr. Biol.* 10, R436 (2000).
5. F. Slack, G. Ruvkun, *Annu. Rev. Genet.* 31, 611 (1997).
6. A. E. Pasquinelli et al., *Nature* 408, 86 (2000).
7. S. M. Elbashir et al., *Nature* 411, 494 (2001).
- 10 8. S. M. Elbashir, W. Lendeckel, T. Tuschl, *Genes & Dev.* 15, 188 (2001).
9. A. J. Hamilton, D. C. Baulcombe, *Science* 286, 950 (1999).
10. S. M. Hammond, E. Bernstein, D. Beach, G. J. Hannon, *Nature* 404, 293 (2000).
- 15 11. P. D. Zamore, T. Tuschl, P. A. Sharp, D. P. Bartel, *Cell* 101, 25 (2000).
12. G. Hutvagner, J. McLachlan, É. Bálint, T. Tuschl, P. D. Zamore, *Science* 93, 834 (2001).
13. A. Grishok et al., *Cell* 106, 23 (2001).
- 20 14. Cloning of 19- to 24-nt RNAs from *D. melanogaster* 0-2 h embryo lysate was performed as described (8). For cloning of HeLa miRNAs, 1 mg of HeLa total RNA was separated on a 15% denaturing polyacrylamide gel and RNA of 19- to 25-nt size was recovered. A 5' phosphorylated 3' adapter oligonucleotide (5' pUUU-aaccgcgaattccagx: uppercase, RNA; lowercase, DNA; p, phosphate; x, 4-hydroxymethylbenzyl; SEQ ID NO:54) and a 5' adapter oligonucleotide (5' acggaattcctcactAAA: uppercase, RNA; lowercase, DNA; SEQ ID NO:55) were ligated to the short HeLa cell RNAs. RT/PCR was performed with 3' primer (5' GACTAGCTGGAATTCGCGGTAAA; SEQ ID NO:56) and 5' primer (5' CAGCCAACGGAATTCCTCACTAAA; SEQ ID NO:57), and followed by concatamerization after Eco RI digestion and T4 DNA
- 30



ligation (8). After ligation of concatamers into pCR2.1 TOPO vectors, about 100 clones were selected and subjected to sequencing.

15. I. Schneider, *J Embryol Exp Morphol* 27, 353 (1972).
- 5 16. R. Feinbaum, V. Ambros, *Dev. Biol.* 210, 87 (1999).
17. S. M. Hammond, S. Boettcher, A. A. Caudy, R. Kobayashi, G. J. Hannon, *Science* 293, 1146 (2001).
18. A. A. Aravin et al., *Curr. Biol.* 11, 1017 (2001).
19. H. Tabara et al., *Cell* 99, 123 (1999).
- 10 20. M. Fagard, S. Boutet, J. B. Morel, C. Bellini, H. Vaucheret, *Proc. Natl. Acad. Sci. USA* 97, 11650 (2000).
21. C. Catalanotto, G. Azzalin, G. Macino, C. Cogoni, *Nature* 404, 245 (2000).
22. S. R. Eddy, *Curr. Opin. Genet. Dev.* 9, 695 (1999).
- 15 23. J. Cavaille et al., *Proc. Natl. Acad. Sci. USA* 97, 14311 (2000).
24. A. Hüttenhofer et al., *EMBO J.* 20, 2943 (2001).
25. L. Argaman et al., *Curr. Biol.* 11, 941 (2001).
26. K. M. Wassarman, F. Repoila, C. Rosenow, G. Storz, S. Gottesman, *Genes & Dev.* 15, 1637 (2001).
- 20 27. Supplementary Web material is available on Science Online at [www.sciencemag.org/cgi/content/full/xxx](http://www.sciencemag.org/cgi/content/full/xxx)
28. D. H. Mathews, J. Sabina, M. Zuker, D. H. Turner, *J. Mol. Biol.* 288, 911 (1999).
29. E. Bernstein, A. A. Caudy, S. M. Hammond, G. J. Hannon, *Nature* 25 409, 363 (2001).
30. Graham, F.L. and van der Eb, A.J., (1973), *Virol.* 52, 456.
31. McCutchan, J.H. and Pagano, J.S., (1968), *J. Natl. Cancer Inst.* 41, 351.
32. Chu, G. et al., (1987), *Nucl. Acids Res.* 15, 1311.
- 30 33. Fraley, R. et al., (1980), *J. Biol. Chem.* 255, 10431.
34. Capecchi, M.R., (1980), *Cell* 22, 479.
35. Felgner, P.L. et al., (1987), *Proc. Natl. Acad. Sci USA* 84, 7413.

36. Lau N.C., Lim L.P., Weinstein E.G., Bartel D.P., (2001), *Science* 294, 858-862.
37. Lee R.C., Ambros V., (2001), *Science* 294, 862-864.
38. Ambros V., (2001), *Cell* 107, 823-826.
- 5 39. Ambros V., Horvitz H.R., (1984), *Science* 226, 409-416.
40. Wightman B., Ha I., Ruvkun G., (1993), *Cell* 75, 855-862.
41. Rougvie A.E., (2001), *Nat. Rev. Genet.* 2, 690-701.
42. Ketting R.F., Fischer S.E., Bernstein E., Sijen T., Hannon G.J.,  
Plasterk R.H., (2001), *Genes & Dev.* 15, 2654-2659.
- 10 43. Hallam S.J., Jin Y., (1998), *Nature* 395, 78-82.
44. Gauwerky C.E., Huebner K., Isobe M., Nowell P.C., Croce C.M.,  
(1989), *Proc. Natl. Acad. Sci. USA* 86, 8867-8871.
45. P. Chomczynski, N. Sacchi, *Anal Biochem* 162, 156, (1987).
46. Mourelatos Z., Dostie J., Paushkin S., Sharma A., Charroux B., Abel  
15 L., J.R., Mann M., Dreyfuss G., (2002), *Genes & Dev.*, in press.
47. Tam W., (2001), *Gene* 274, 157-167.

**Table 1**

D. melanogaster miRNAs. The sequences given represent the most abundant, and typically longest miRNA sequence identified by cloning; miRNAs frequently vary in length by one or two nucleotides at their 3' termini. From 222 short RNAs sequenced, 69 (31%) corresponded to miRNAs, 103 (46%) to already characterized functional RNAs (rRNA, 7SL RNA, tRNAs), 30 (14%) to transposon RNA fragments, and 20 (10%) sequences with no database entry. The frequency (freq.) for cloning a particular miRNA relative to all identified miRNAs is indicated in percent.

Results of Northern blotting of total RNA isolated from staged populations of D. melanogaster are summarized. E, embryo; L, larval stage; P, pupae; A, adult; S2, Schneider-2 cells. The strength of the signal within each blot is represented from strongest (+ + +) to undetected (-). let-7 stRNA was probed as control. Genbank accession numbers and homologs of miRNAs identified by database searching in other species are provided as supplementary material.

miRNA	sequence (5' to 3')	freq. (%)	E 0-3 h	E 0-6 h	L1+ L2	L3	P	A	S2
miR-1	UGGAAUGUAAAAGAAGUAUGGAG (SEQ ID NO:58)	32	+	+	++ +	++ +	++	++ +	-
miR-2a*	UAUCACAGCCAGCUUUGAUGAGC (SEQ ID NO:59)	3							
miR-2b*	UAUCACAGCCAGCUUUGAGGAGC (SEQ ID NO:60)	3	++	++	++ +	++ +	++	+	++ +
miR-3	UCACUGGGCAAAGUGUGUCUCA#	9	+++	+++	-	-	-	-	-
miR-4	AUAAAGCUAGACAACCAUUGA (SEQ ID NO:62)	6	+++	+++	-	-	-	-	-
miR-5	AAAGGAACGAUCGUUGUGAUAUG (SEQ ID NO:63)	1	+++	+++	+/-	+/-	-	-	-
miR-6	UAUCACAGUGGCUGUUCUUUUU (SEQ ID NO:64)	13	+++	+++	+/-	+/-	-	-	-
miR-7	UGGAAGACUAGUGAUUUUGUUGU (SEQ ID NO:65)	4	+++	++	+/-	+/-	+/-	+/-	+/-
miR-8	UAAUACUGUCAGGUAAAAGAUGUC (SEQ ID NO:66)	3	+/-	+/-	++ +	++ +	+	++ +	-

- 27 -

miR-9	UCUUUGGUUAUCUAGCUGUAUGA (SEQ ID NO:67)	7	+++	++	++	++	++	+/-	-
miR-10	ACCCUGUAGAUCGAAUUUGU (SEQ ID NO:68)	1	+	+	++	++	+/-	+	-
miR-11	CAUCACAGUCUGAGUUCUUGC (SEQ ID NO:69)	7	+++	+++	++	++	++	+	-
miR-12	UGAGUAUUACAUCAGGUACUGGU (SEQ ID NO:70)	7	+	+	++	++	+	++	+/-
5 miR-13a*	UAUCACAGCCAUUUUGACGAGU (SEQ ID NO:71)	1	+++	+++	++	++	+	++	++
miR-13b*	UAUCACAGCCAUUUUGAUGAGU (SEQ ID NO:72)	0							
miR-14	UCAGUCUUUUUCUCUCCUA (SEQ ID NO:73)	1	-	-	-	-	-	-	-
let-7	UGAGGUAGUAGGUUGUAUAGUU (SEQ ID NO:74)	0	-	-	-	-	++	++	-

10 # = (SEQ ID NO:61)

\*Similar miRNA sequences are difficult to distinguish by Northern blotting because of potential cross-hybridization of probes.

Table 2

Human miRNAs. From 220 short RNAs sequenced, 100 (45%) corresponded to miRNAs, 53 (24%) to already characterized functional RNAs (rRNA, snRNAs, tRNAs), and 67 (30%) sequences with no database entry. Results of Northern blotting of total RNA isolated from different vertebrate species and S2 cells are indicated. For legend, see Table 1.

miRNA	sequence (5' to 3')	freq. (%)	HeLa cells	mouse kidney	adult fish	frog ovary	S2
10 let-7a*	UGAGGUAGUAGGUUGUAUAGUU#	10	+++	+++	+++	-	-
let-7b*	UGAGGUAGUAGGUUGUGUGGUU (SEQ ID NO:76)	13					
let-7c*	UGAGGUAGUAGGUUGUAUGGUU (SEQ ID NO:77)	3					
let-7d*	AGAGGUAGUAGGUUGCAUAGU (SEQ ID NO:78)	2	+++	+++	+++	-	-
let-7e*	UGAGGUAGGAGGUUGUAUAGU (SEQ ID NO:79)	2	+++	+++	+++	-	-
let-7f*	UGAGGUAGUAGAUUGUAUAGUU (SEQ ID NO:80)	1					
15 miR-15	UAGCAGCACAAUAAUGGUUUGUG (SEQ ID NO:81)	3	+++	++	+	+/-	-
miR-16	UAGCAGCACGUAAAUAUUGGCG (SEQ ID NO:82)	10	+++	+	+/-	+/-	-
miR-17	ACUGCAGUGAAGGCACUUGU (SEQ ID NO:83)	1	+++	-	-	-	-
miR-18	UAAGGUGCAUCUAGUGCAGUA (SEQ ID NO:84)	2	+++	-	-	-	-
miR-19a*	UGUGCAAUUCUAUGCAAACUGA (SEQ ID NO:85)	1	+++	-	+/-	-	-
20 miR-19b*	UGUGCAAUCCAUGCAAACUGA (SEQ ID NO:86)	3					
miR-20	UAAAGUGCUUAUAGUGCAGGUA (SEQ ID NO:87)	4	+++	-	+	-	-
miR-21	UAGCUUAUCAGACUGAUGUUGA (SEQ ID NO:88)	10	+++	+	++	-	-
miR-22	AAGCUGCCAGUUGAAGAACUGU (SEQ ID NO:89)	10	+++	+++	+	+/-	-
miR-23	AUCACAUUGCCAGGGAUUCC (SEQ ID NO:90)	2	+++	+++	+++	+	-

- 29 -

	miR-24	UGGCUCAGUUCAGCAGGAACAG (SEQ ID NO:91)	4	++	+++	++	-	-
	miR-25	CAUUGCACUUGUCUCGGUCUGA (SEQ ID NO:92)	3	+++	+	++	-	-
	miR-26a*	UUCAAGUAAUCCAGGAUAGGCU (SEQ ID NO:93)	2	+	++	+++	-	-
	miR-26b*	UUCAAGUAAUUCAGGAUAGGUU (SEQ ID NO:94)	1					-
5	miR-27	UUCACAGUGGCUAAGUUCGCGU (SEQ ID NO:95)	2	+++	+++	++	-	-
	miR-28	AAGGAGCUCACAGUCUAUUGAG (SEQ ID NO:96)	2	+++	+++	-	-	-
	miR-29	CUAGCACCAUCUGAAAUCGGUU (SEQ ID NO:97)	2	+	+++	+/-	-	-
	miR-30	CUUCAGUCGGAUGUUUCAGC (SEQ ID NO:98)	2	+++	+++	+++	-	-
	miR-31	GGCAAGAUGCUGGCAUAGCUG (SEQ ID NO:99)	2	+++	-	-	-	-
10	miR-32	UAUUGCACAUACUAAGUUGC (SEQ ID NO:100)	1	-	-	-	-	-
	miR-33	GUGCAUUGUAGUUGCAUUG (SEQ ID NO:101)	1	-	-	-	-	-
	miR-1	UGGAAUGUAAAGAAGUAUGGAG (SEQ ID NO:102)	0	-	-	+	-	-
	miR-7	UGGAAGACUAGUGAUUUUGUUGU (SEQ ID NO:103)	0	+	-	+/-	-	+/-
	miR-9	UCUUUGGUUAUCUAGCUGUAUGA (SEQ ID NO:104)	0	-	-	-	-	-
15	miR-10	ACCCUGUAGAUCCGAAUUUGU (SEQ ID NO:105)	0	-	+	-	-	-

# = (SEQ ID NO:75)

\*Similar miRNA sequences are difficult to distinguish by Northern blotting because of potential cross-hybridization of probes.

- 30 -

**Table 3**

Mouse miRNAs. The sequences indicated represent the longest miRNA sequences identified by cloning. The 3'-terminus of miRNAs is often truncated by one or two nucleotides. miRNAs that are more than 85% identical in sequence (i.e. share 18 out of 21 nucleotides) or contain 1- or 2-nucleotide internal deletions are referred to by the same gene number followed by a lowercase letter. Minor sequence variations between related miRNAs are generally found near the ends of the miRNA sequence and are thought to not compromise target RNA recognition. Minor sequence variations may also represent A to G and C to U changes, which are accommodated as G-U wobble base pairs during target recognition. miRNAs with the suffix -s or -as indicate RNAs derived from either the 5'-half or the 3'-half of a miRNA precursor. Mouse brains were dissected into midbrain, mb, cortex, cx, cerebellum, cb. The tissues analyzed were heart, ht; liver, lv; small intestine, si; colon, co; cortex, ct; cerebellum, cb; midbrain, mb.

miRNA	sequence (5' to 3')	Number of clones							
		ht	lv	sp	si	co	cx	cb	mb
let-7a	UGAGGUAGUAGGUUGUAUAGUU (SEQ ID NO:106)	3				1	1		7
let-7b	UGAGGUAGUAGGUUGUGUGUU (SEQ ID NO:107)	1	1					2	5
let-7c	UGAGGUAGUAGGUUGUAUGUU (SEQ ID NO:108)	2					2	5	19
let-7d	AGAGGUAGUAGGUUGCAUAGU (SEQ ID NO:109)	2				2	2		2
let-7e	UGAGGUAGGAGGUUGUAUAGU (SEQ ID NO:110)			1					2
let-7f	UGAGGUAGUAGAUUGUAUAGUU (SEQ ID NO:111)			2				3	3
let-7g	UGAGGUAGUAGUUUGUACAGUA (SEQ ID NO:112)						1	1	2
let-7h	UGAGGUAGUAGUGUGUACAGUU (SEQ ID NO:113)						1	1	

	let-7i	UGAGGUAGUAGUUUGUCU (SEQ ID NO:114)				1	1		
	miR-1b	UGGAAUGUAAAGAAGUAUGUAA (SEQ ID NO:115)	4	2					1
	miR-1c	UGGAAUGUAAAGAAGUAUGUAC (SEQ ID NO:116)	7						
	miR-1d	UGGAAUGUAAAGAAGUAUGUAUU (SEQ ID NO:117)	16						1
5	miR-9	UCUUUGGUUAUCUAGCUGUAUGA (SEQ ID NO:118)				3	4		4
	miR-15a	UAGCAGCACAUAAUGGUUUGUG (SEQ ID NO:119)	1						2
	miR-15b	UAGCAGCACAUCAUGGUUACA (SEQ ID NO:120)	1						
	miR-16	UAGCAGCACGUAAAUUUGGCG (SEQ ID NO:121)	1			1	2	1	2
	miR-18	UAAGGUGCAUCUAGUGCAGUA (SEQ ID NO:122)				1			
10	miR-19b	UGUGCAAUCCAUGCAAACUGA (SEQ ID NO:123)				1			
	miR-20	UAAAGUGCUUAUAGUGCAGGUAG (SEQ ID NO:124)				1			
	miR-21	UAGCUUAUCAGACUGAUGUUGA (SEQ ID NO:125)	1			1	2	1	
	miR-22	AAGCUGCCAGUUGAAGAACUGU (SEQ ID NO:126)	2	1		1		1	2
	miR-23a	AUCACAUUGCCAGGGAUUUC (SEQ ID NO:127)	1						
15	miR-23b	AUCACAUUGCCAGGGAUUACCAC (SEQ ID NO:128)				1			
	miR-24	UGGCUCAGUUCAGCAGGAACAG (SEQ ID NO:129)	1			1	1		1
	miR-26a	UUCAAGUAAUCCAGGAUAGGCU (SEQ ID NO:130)						3	2
	miR-26b	UUCAAGUAAUCCAGGAUAGGUU (SEQ ID NO:131)		2			4	1	
	miR-27a	UUCACAGUGGCUAAGUCCGCU (SEQ ID NO:132)	1			2	1	1	2
20	miR-27b	UUCACAGUGGCUAAGUUCUG (SEQ ID NO:133)							1
	miR-29a	CUAGCACCAUCUGAAAUCGGUU (SEQ ID NO:134)	1			1		1	
	miR-29b/miR-102	UAGCACCAUUGAAAUCAGUGUU (SEQ ID NO:135)	1			1	5		3
	miR-29c/	UAGCACCAUUGAAAUCGGUUA (SEQ ID NO:136)	1			3			1



	miR-30a-s/miR-97	UGUAAACAUCCUCGACUGGAAGC (SEQ ID NO:137)	1		1		1
	miR-30a-as <sup>a</sup>	CUUUCAGUCGGAUGUUUGCAGC (SEQ ID NO:138)					1
	miR-30b	UGUAAACAUCCUACACUCAGC (SEQ ID NO:139)	1				2
	miR-30c	UGUAAACAUCCUACACUCUCAGC (SEQ ID NO:140)	2			1	1
5	miR-30d	UGUAAACAUCCCGACUGGAAG (SEQ ID NO:141)		1			
	miR-99a/miR-99	ACCCGUAGAUCGGAUCUUGU (SEQ ID NO:142)				1	
	miR-99b	CACCCGUAGAACCGACCUUGCG (SEQ ID NO:143)					1
	miR-101	UACAGUACUGUGAUAAACUGA (SEQ ID NO:144)				2	1 1
	miR-122a	UGGAGUGUGACAAUGGUGUUUGU (SEQ ID NO:145)		3			
10	miR-122b	UGGAGUGUGACAAUGGUGUUUGA (SEQ ID NO:146)		11			
	miR-122a,b	UGGAGUGUGACAAUGGUGUUUG (SEQ ID NO:147)		23			
	miR-123	CAUUAUUACUUUUGGUACGCG (SEQ ID NO:148)	1	2			
	miR-124a <sup>b</sup>	UUAAGGCACGCGG-UGAAUGCCA (SEQ ID NO:149)		1		37	41 24
	miR-124b	UUAAGGCACGCGGGUGAAUGC (SEQ ID NO:150)				1	3
15	miR-125a	UCCUGAGACCCUUUAACCGUG (SEQ ID NO:151)				1	1
	miR-125b	UCCUGAGACCCU--AACUUGUGA (SEQ ID NO:152)				1	
	miR-126	UCGUACCGUGAGUAAUAAUGC (SEQ ID NO:153)	4				1
	miR-127	UCGGAUCCGUCUGAGCUUGGCU (SEQ ID NO:154)					1
	miR-128	UCACAGUGAACCGGUCUCUUUU (SEQ ID NO:155)				2	2 2
20	miR-129	CUUUUUUCGGUCUGGGCUUGC (SEQ ID NO:156)					1
	miR-130	CAGUGCAAUGUAAAAGGGC (SEQ ID NO:157)					1
	miR-131	UAAAGCUAGAUACCGAAAGU (SEQ ID NO:158)				1	1 1
	miR-132	UAACAGUCUACAGCCAUGGUCGU (SEQ ID NO:159)					1

	miR-133	UUGGUCCCCUUCACCAGCUGU (SEQ ID NO:160)	4			1	
	miR-134	UGUGACUGGUUGACCAGAGGGA (SEQ ID NO:161)				1	
	miR-135	UAUGGCUUUUUAUUCCUAUGUGAA (SEQ ID NO:162)				1	
	miR-136	ACUCCAUUUGUUUGAUGAUGGA (SEQ ID NO:163)				1	
5	miR-137	UAUUGCUUAAGAAUCGCGUAG (SEQ ID NO:164)				1	1
	miR-138	AGCUGGUGUUGUGAAUC (SEQ ID NO:165)				1	
	miR-139	UCUACAGUGCACGUGUCU (SEQ ID NO:166)				1	1
	miR-140	AGUGGUUUACCCUAUGGUAG (SEQ ID NO:167)				1	
	miR-141	AACACUGUCUGGUAAGAUGG (SEQ ID NO:168)		1	1		1
10	miR-142-s	CAUAAAGUAGAAAGCACUAC (SEQ ID NO:169)				1	1
	miR-142-as <sup>b</sup>	UGUAGUGUUCCUACUUUAUGG (SEQ ID NO:170)		1	1		6
	miR-143	UGAGAUGAAGCACUGUAGCUA (SEQ ID NO:171)	3		7		2 1
	miR-144	UACAGUAUAGAUGAUGUACUAG (SEQ ID NO:172)	2			1	
	miR-145	GUCCAGUUUCCAGGAAUCCCUU (SEQ ID NO:173)	1				
15	miR-146	UGAGAACUGAAUCCAUGGGUUU (SEQ ID NO:174)	1				
	miR-147	GUGUGUGGAAAUGCUUCUGCC (SEQ ID NO:175)			1		
	miR-148	UCAGUGCACUACAGAACUUUGU (SEQ ID NO:176)			1		
	miR-149	UCUGGCUCGGUGUCUUCACUCC (SEQ ID NO:177)	1				
	miR-150	UCUCCCAACCCUUGUACCAGUGU (SEQ ID NO:178)				1	
20	miR-151	CUAGACUGAGGCUCUUGAGGU (SEQ ID NO:179)				1	
	miR-152	UCAGUGCAUGACAGAACUUGG (SEQ ID NO:180)				1	
	miR-153	UUGCAUAGUCACAAAAGUGA (SEQ ID NO:181)					1
	miR-154	UAGGUUAUCCGUGUUGCCUUCG (SEQ ID NO:182)					1

- 34 -

miR-155

UUA AUGCUAAUUGUGAUAGGGG  
(SEQ ID NO:183)

1

5 The originally described miR-30 was renamed to miR-30a-as in order to distinguish it from the miRNA derived from the opposite strand of the precursor encoded by the mir-30a gene. miR-30a-s is equivalent to miR-97 [46].

<sup>b</sup>A 1-nt length heterogeneity is found on both 5' and 3' end. The 22-nt miR sequence is shown, but only 21-nt miRNAs were cloned.

10

**Table 4**

Mouse and human miRNAs. The sequences indicated represent the longest miRNA sequences identified by cloning. The 3' terminus of miRNAs is often truncated by one or two nucleotides. miRNAs that are more than 85% identical in sequence (i.e. share 18 out of 21 nucleotides) or contain 1- or 2-nucleotide internal deletions are referred to by the same gene number followed by a lowercase letter. Minor sequence variations between related miRNAs are generally found near the ends of the miRNA sequence and are thought to not compromise target RNA recognition. Minor sequence variations may also represent A to G and C to U changes; which are accommodated as G-U wobble base pairs during target recognition. Mouse brains were dissected into midbrain, mb, cortex, cx, cerebellum, cb. The tissues analyzed were lung, ln; liver, lv; spleen, sp; kidney, kd; skin, sk; testis, ts; ovary, ov; thymus, thy; eye, ey; cortex, ct; cerebellum, cb; midbrain, mb. The human osteosarcoma cells SAOS-2 cells contained an inducible p53 gene (p53-, uninduced p53; p53+, induced p53); the differences in miRNAs identified from induced and uninduced SAOS cells were not statistically significant.

number of clones

miRNA	Sequence (5' to 3')	mouse tissues										human SAOS-2 cells			
		ln	lv	sp	kd	sk	ts	ov	thy	ey	p53-	p53+			
miR-C1	AACAUUC AACCGCUCGUGGUGAGU	1				1							2		(SEQ ID NO.184)
miR-C2	UUUGGCAAUGGUAAGAACUCACA												1		(SEQ ID NO.185)
miR-C3	UAUGGCACUGGUAAGAAUUCACUG												1		(SEQ ID NO.186)
miR-C4	CUUUUGCGGUCUGGGCUTUGUU							1					1	1	(SEQ ID NO.187)
miR-C5	UGGACGGAGAACUGAAUAAGGGU												2		(SEQ ID NO.188)
miR-C6	UGGAGAGAAAGGCAGUUC												1		(SEQ ID NO.189)
miR-C7	CAAAGAAUUCUCCUUUGGGCUU												1	1	(SEQ ID NO.190)
miR-C8	UCGUGUCUUGUGUUGCAGCCGG								1						(SEQ ID NO.191)
miR-C9	UAACACUGUCUGGUAACGAUG								1						(SEQ ID NO.192)
miR-C10	CAUCCCUUGCAUGGUGGAGGGU								1						(SEQ ID NO.193)
miR-C11	GUGCCUACUGAGCUGACAUCAGU								1						(SEQ ID NO.194)
miR-C12	UGAUAUGUUUGAUAUAUUAAGU								2						(SEQ ID NO.195)
miR-C13	CAACGGAAUCCCAAAAGCAGCU							2	1						(SEQ ID NO.196)
miR-C14	CUGACCUAUGAAUUGACA							2	1						(SEQ ID NO.197)

miR-C15	UACCACAGGGUAGAACCCAGGA	1		(SEQ ID NO.198)	
miR-C16	AACUGGCCUACAAGUCCAG	1		(SEQ ID NO.199)	
miR-C17	UGUAAACAGCAACUCCAUGUGGA	1		(SEQ ID NO.200)	
miR-C18	UAGCAGCACAGAAAUAUUGGC	2	1	(SEQ ID NO.201)	
5	miR-C19	UAGGUAGUUUCAUGUUGUUGG	1	1	(SEQ ID NO.202)
miR-C20	UUCACCACCUUCCACCCAGC	1	1	(SEQ ID NO.203)	
miR-C21	GGUCCAGAGGGAGAUAGG	1	1	(SEQ ID NO.204)	
miR-C22	CCCAGUGUUCAGACUACCUGUU	1	1	(SEQ ID NO.205)	
miR-C23	UAAUACUGCCUGGUAUUGAC	2	1	(SEQ ID NO.206)	
10	miR-C24	UACUCAGUAAGGCAUUGUUCU	1	1	(SEQ ID NO.207)
miR-C25	AGAGUAUAGCGCAUGGGAAGA	1	1	(SEQ ID NO.208)	
miR-C26	UGAAUUGUUUAGGACCACUAG	2	1	(SEQ ID NO.209)	
miR-C27	UUCUUUUGUCAUCCUAUGCCUG	1	1	(SEQ ID NO.210)	
miR-C28	UCCUUAUCCACCGGAGUCUG	1	1	(SEQ ID NO.211)	
15	miR-C29	GUGAAUUGUUUAGGACCACUAGA	2	1	(SEQ ID NO.212)
miR-C30	UGGAAUGUAAGGAGUGUGUGG	2	1	(SEQ ID NO.213)	
miR-C31	UACAGUAGUCUGCAUUGGUU	1	1	(SEQ ID NO.214)	
miR-C32	CCUUGUAGAACCGAAUUGUGU	1	1	(SEQ ID NO.215)	
miR-C33	AACCCGUAGAUCCGAACUUGUGAA	1	1	(SEQ ID NO.216)	
20	miR-C34	GCUUCUCCUGGCUCUCCUCCUC	1	1	(SEQ ID NO.217)

Table 5

*D. melanogaster* miRNA sequences and genomic location. The sequences given represent the most abundant, and typically longest miRNA sequences identified by cloning. It was frequently observed that miRNAs vary in length by one or two nucleotides at their 3'-terminus. From 222 short RNAs sequenced, 69 (31%) corresponded to miRNAs, 103 (46%) to already characterized functional RNAs (rRNA, 7SL RNA, tRNAs), 30 (14%) to transposon RNA fragments, and 20 (10%) sequences with no database entry. RNA sequences with a 5'-guanosine are likely to be underrepresented due to the cloning procedure (8). miRNA homologs found in other species are indicated. Chromosomal location (chr.) and GenBank accession numbers (acc. nb.) are indicated. No ESTs matching miR-1 to miR-14 were detectable by database searching.

miRNA	sequence (5' to 3')	chr., acc. nb.	remarks
miR-1	UGGAAUGUAAAGAAGUAUGGAG (SEQ ID NO: 58)	2L, AE003667	homologs: <i>C. briggsae</i> , G20U, AC87074; <i>C. elegans</i> G20U, U97405; mouse, G20U, G22U, AC020867; human, chr. 20, G20U, G22U, AL449263; ESTs: zebrafish, G20U, G22U, BF157-601; cow, G20U, G22U, BE722-224; human, G20U, G22U, AI220268
miR-2a	UAUCACAGCCAGCUUUGAUGAGC (SEQ ID NO: 59)	2L, AE003663	2 precursor variants clustered with a copy of <i>mir-2b</i>
miR-2b	UAUCACAGCCAGCUUUGAGGAGC (SEQ ID NO: 60)	2L, AE003620 2L, AE003663	2 precursor variants
miR-3	UCACUGGGCAAAGUGUGUCUCA (SEQ ID NO: 61)	2R, AE003795	In cluster <i>mir-3</i> to <i>mir-6</i>
miR-4	AUAAAGCUAGACAACCAUUGA (SEQ ID NO: 62)	2R, AE003795	in cluster <i>mir-3</i> to <i>mir-6</i>

- 39 -

	miR-5	AAAGGAACGAUCGUUGUGAUUUG (SEQ ID NO:63)	2R, AE003795	in cluster <i>mir-3</i> to <i>mir-6</i>
	miR-6	UAUCACAGUGGCUGUUCUUUUU (SEQ ID NO:64)	2R, AE003795	in cluster <i>mir-3</i> to <i>mir-6</i> with 3 variants
5	miR-7	UGGAAGACTAGUGAUUUUGUUGU (SEQ ID NO:65)	2R, AE003791	homologs: human, chr. 19 AC006537, EST BF373391; mouse chr. 17 AC026385, EST AA881786
	miR-8	UAAUACUGUCAGGUAAAAGAUGUC (SEQ ID NO:66)	2R, AE003805	
10	miR-9	UCUUUGGUUAUCUAGCUGUAUGA (SEQ ID NO:67)	3L, AE003516	homologs: mouse, chr. 19, AF155142; human, chr. 5, AC026701, chr. 15, AC005316
	miR-10	ACCCUGUAGAUCCGAAUUUGU (SEQ ID NO:68)	AE001574	homologs: mouse, chr 11, AC011194; human, chr. 17, AF287967
	miR-11	CAUCACAGUCUGAGUUCUUUGC (SEQ ID NO:69)	3R, AE003735	intronic location
15	miR-12	UGAGUAUUACAUCAGGUACUGGU (SEQ ID NO:70)	X, AE003499	intronic location
	miR-13a	UAUCACAGCCAUUUUGACGAGU (SEQ ID NO:71)	3R, AE003708 X, AE003446	<i>mir-13a</i> clustered with <i>mir-13b</i> on chr. 3R
20	miR-13b	UAUCACAGCCAUUUUGAUGAGU (SEQ ID NO:72)	3R, AE003708	<i>mir-13a</i> clustered with <i>mir-13b</i> on chr. 3R
	miR-14	UCAGUCUUUUUCUCUCUCCUA (SEQ ID NO:73)	2R, AE003833	no signal by Northern analysis

---



Table 6

Human miRNA sequences and genomic location. From 220 short RNAs sequenced, 100 (45%) corresponded to miRNAs, 53 (24%) to already characterized functional RNAs (rRNA, snRNAs, tRNAs), and 67 (30%) sequences with no database entry. For legend, see Table 1.

miRNA	sequence (5' to 3')	chr. or EST, acc. nb.	remarks*
10 let-7a	UGAGGUAGUAGGUUGUAUAGUU (SEQ ID NO:75)	9, AC007924, 11, AP001359, 17, AC087784, 22, AL049853	sequences of chr 9 and 17 identical and clustered with <i>let-7f</i> , homologs: <i>C. elegans</i> , AF274345; <i>C. briggsae</i> , AF210771, <i>D.</i> <i>melanogaster</i> , AE003659
let-7b	UGAGGUAGUAGGUUGUGUGGUU (SEQ ID NO:76)	22, AL049853†, ESTs, AI382133, AW028822	homologs: mouse, EST AI481799; rat, EST, BE120662
let-7c	UGAGGUAGUAGGUUGUAUGGUU (SEQ ID NO:77)	21, AP001667	Homologs: mouse, EST, AA575575
15 let-7d	AGAGGUAGUAGGUUGCAUAGU (SEQ ID NO:78)	17, AC087784, 9, AC007924	Identical precursor sequences
let-7e	UGAGGUAGGAGGUUGUAUAGU (SEQ ID NO:79)	19, AC018755	
20 let-7f	UGAGGUAGUAGAUGUAUAGUU (SEQ ID NO:80)	9, AC007924, 17, AC087784, X, AL592046	sequences of chr 9 and 17 identical and clustered with <i>let-7a</i>
miR-15	UAGCAGCACAUAAUGGUUUGUG (SEQ ID NO:81)	13, AC069475	in cluster with <i>mir-16</i> homolog
miR-16	UAGCAGCACGUAAUAUUGGCG (SEQ ID NO:82)	13, AC069475	in cluster with <i>mir-15</i> homolog

- 41 -

	miR-17	ACUGCAGUGAAGGCACUUGU (SEQ ID NO:83)	13, AL138714	in cluster with <i>mir-17</i> to <i>mir-20</i>
	miR-18	UAAGGUGCAUCUAGUGCAGAU (SEQ ID NO:84)	13, AL138714	in cluster with <i>mir-17</i> to <i>mir-20</i>
5	miR-19a	UGUGCAAUUCUAUGCAAACUG A (SEQ ID NO:85)	13, AL138714	in cluster with <i>mir-17</i> to <i>mir-20</i>
	miR-19b	UGUGCAAUCCAUGCAAACUG A (SEQ ID NO:86)	13, AL138714, X, AC002407	in cluster with <i>mir-17</i> to <i>mir-20</i>
10	miR-20	UAAAGUGCUUAUAGUGCAGGUA (SEQ ID NO:87)	13, AL138714	in cluster with <i>mir-17</i> to <i>mir-20</i>
	miR-21	UAGCUUAUCAGACTUGAUGUUGA (SEQ ID NO:88)	17, AC004686, EST, BF326048	homologs: mouse, EST, AA209594
	miR-22	AAGCUGCCAGUUGAAGAACUGU (SEQ ID NO:89)	ESTs, AW961681†, AA456477, AI752503, BF030303, HS1242049	human ESTs highly similar; homologs: mouse, ESTs, e.g. AA823029; rat, ESTs, e.g. BF543690
15	miR-23	AUCACAUUGCCAGGGAUUCU (SEQ ID NO:90)	19, AC020916	homologs: mouse, EST, AW124037;rat, EST, BF402515
	miR-24	UGGCUCAGUUCAGCAGGAACAG (SEQ ID NO:91)	9, AF043896, 19, AC020916	homologs: mouse, ESTs, AA111466, AI286629; pig, EST, BE030976
20	miR-25	CAUUGCACUUGUCUCGGUCUGA (SEQ ID NO:92)	7, AC073842, EST, BE077684	human chr 7 and EST identical; highly similar precursors in mouse ESTs (e.g. AI595464); fish precursor different STS: G46757
	miR-26a	UUCAAGUAAUCCAGGAUAGGCU (SEQ ID NO:93)	3, AP000497	

- 42 -

	miR-26b	UUCAAGUAAUUCAGGAUAGGUU (SEQ ID NO:94)	2, AC021016	
	miR-27	UUCACAGUGGCUAAGUCCGCU (SEQ ID NO:95)	19, AC20916	U22C mutation in human genomic sequence
5	miR-28	AAGGAGCUCACAGUCUAUUGAG (SEQ ID NO:96)	3, AC063932	
	miR-29	CUAGCACCAUCUGAAAUCGGUU (SEQ ID NO:97)	7, AF017104	
	miR-30	CUUUCAGUCGGAUGUUUGCAGC (SEQ ID NO:98)	6, AL035467	
10	miR-31	GGCAAGAUGCUGGCAUAGCUG (SEQ ID NO:99)	9, AL353732	
	miR-32	UAUUGCACAUUACUAAGUUGC (SEQ ID NO:100)	9, AL354797	not detected by Northern blotting
15	miR-33	GUGCAUUGUAGUUGCAUUG (SEQ ID NO:101)	22, Z99716	not detected by Northern blotting

\*If several ESTs were retrieved for one organism in the database, only those with different precursor sequences are listed.

20 †precursor structure shown in Fig. 4.

### Claims

1. Isolated nucleic acid molecule comprising
  - (a) a nucleotide sequence as shown in Table 1, Table 2, Table 3 or Table 4 or a precursor thereof as shown in Figure 3, Figure 4 or Figure 7.
  - (b) a nucleotide sequence which is the complement of (a),
  - (c) a nucleotide sequence which has an identity of at least 80% to a sequence of (a) or (b) and/or
  - (d) a nucleotide sequence which hybridizes under stringent conditions to a sequence of (a), (b) and/or (c).
2. The nucleic acid molecule of claim 1, wherein the identity of sequence (c) is at least 90%.
3. The nucleic acid molecule of claim 1, wherein the identity of sequence (c) is at least 95%.
4. The nucleic acid molecule of any one of claims 1-3, which is selected from miR 1-14 as shown in Table 1 or miR 15-33 as shown in Table 2 or miR 1-155 as shown in Table 3 or miR-C1-34 as shown in Table 4 or a complement thereof.
5. The nucleic acid molecule of any one of claims 1-3, which is selected from mir 1-14 as shown in Figure 3 or let 7a-7f or mir 15-33, as shown in Figure 4 or let 7a-i or mir 1-155 or mir-c1-34, as shown in Figure 7 or a complement thereof.

6. The nucleic acid molecule of any one of claims 1-4 which is a miRNA molecule or an analog thereof having a length of from 18-25 nucleotides.
7. The nucleic acid molecule of any one of claims 1-3 or 5, which is a  
5 miRNA precursor molecule having a length of 60-80 nucleotides or a DNA molecule coding therefor.
8. The nucleic acid molecule of any one of claims 1-7, which is single-stranded.
- 10 9. The nucleic acid molecule of any one of claims 1-7, which is at least partially double-stranded.
- 15 10. The nucleic acid molecule of any one of claims 1-9, which is selected from RNA, DNA or nucleic acid analog molecules.
11. The nucleic acid molecule of claim 10, which is a molecule containing at least one modified nucleotide analog.
- 20 12. The nucleic molecule of claim 10 which is a recombinant expression vector.
13. A pharmaceutical composition containinig as an active agent at least one nucleic acid molecule of any one of claims 1-12 and optionally a pharmaceutically acceptable carrier.  
25
14. The composition of claim 13 for diagnostic applications.
15. The composition of claim 13 for therapeutic applications.
- 30 16. The composition of any one of claims 13-15 as a marker or a modulator for developmental or pathogenic processes.

17. The composition of claim 13 as a marker or modulator of developmental disorders, particularly cancer, such a B-cell chronic leukemia.
18. The composition of any one of claims 13-15 as a marker or modulator of gene expression.
19. The composition of claim 18 as a marker or modulator of the expression of a gene, which is at least partially complementary to said nucleic acid molecule.
20. A method of identifying microRNA molecules or precursor molecules thereof comprising ligating 5'- and 3'-adapter molecules to the ends of a size-fractionated RNA population, reverse transcribing said adapter-containing RNA population and characterizing the reverse transcription products.

Fig. 1 A




Fig. 1 B






Fig. 2




Fig. 3






Fig. 5







Fig. 5 (cont.)

brain  
rbmbcx cb ht lg lv co si pc sp kd sm st H




brain  
rbmbcx cb ht lg lv co si pc sp kd sm st H




— miR-L


— miR-S



brain  
rbmbcx cb ht lg lv co si pc sp kd sm st H



brain  
rbmbcx cb ht lg lv co si pc sp kd sm st H




— miR-L


— miR-S

Fig. 5 (cont.)

brain  
rbmbcx cb ht lg lv co si pc sp kd sm st H




brain  
rbmbcx cb ht lg lv co si pc sp kd sm st H




— miR-L  
— miR-S

brain  
rbmbcx cb ht lg lv co si pc sp kd sm st H



brain  
rbmbcx cb ht lg lv co si pc sp kd sm st H



— miR-L  
— miR-S

Fig. 5 (cont.)




Fig. 6

A

<i>C. elegans</i> lin-4	UCCUGAGACCUC--AAG-UGUGA
<i>D. melanogaster</i> miR-125	UCCUGAGACCCU--AAUUGUGA
<i>M. musculus/H. sapiens</i> miR-125b	UCCUGAGACCCU--AAUUGUGA
<i>M. musculus/H. sapiens</i> miR-125a	UCCUGAGACCCUUUAACCUGUGA

B






Fig. 7

name	sequence	structure
let-7a-1	UGAGGUAGUAGGUUGUAVAGUU	<pre>           UG      U      UUAGG ACA C CAC UGGA GAGUAGUAGGUUGUAVAGUU GUC CCCA C GUG AUCCU UUCUGUCAUCAUCAUCAUA CA GGU A           CA - - - - - A - - C           </pre>
let-7a-2	UGAGGUAGUAGGUUGUAVAGUU	<pre>           UU G U UAGAUAUAC AA           AGG GAG UAG AGGUUGUAVAGUU AUC G           UCC UUC AUC UCCGACAUGUCA UAG G           U - G C C - - - - - AG           </pre>
let-7a-3	UGAGGUAGUAGGUUGUAVAGUU	<pre>           U GAGUAGUAGGUUGUAVAGUU U           GGG GAGUAGUAGGUUGUAVAGUU UGGGC \           UCC UUCUGUCAUCAUCAUCAUA GUC CCG C           U UAGGUUAUC U           </pre>
let-7b	UGAGGUAGUAGGUUGUGUGUU	<pre>           U GAGUAGUAGGUUGUGUGUU UG           CGGG GAGUAGUAGGUUGUGUGUU GGCAG \           GUCCC UUCGUAUCAUCAUCAUA CA AG CCCUU A           - - - - - U AAGGCUC GU           </pre>
let-7c	UGAGGUAGUAGGUUGUAVAGUU	<pre>           AUU G U UA G UA AC           GC UCCGG GAG UAG AGGUUGUAVAGUU GA U C \           CG AGGUUC UUC AUC UCCACAUAGUCA UA A G C           - - - - - CU G U -- G GG UC           </pre>
let-7d	AGAGGUAGUAGGUUGCAUAGU	<pre>           A C UUA - - - - - GG           CCUAGGA GAGUAGUAGGUUG AVAGUU GGCAG \           GGAUUCU UUCGUGUCCAGC UAUCA A CCCUU A           - - - - - A UGGAGGAACA UU           </pre>
let-7e	UGAGGUAGGAGGUUGUAVAGU	<pre>           C CU G U GGA - - - - - A           CC GGG GAG UAGGAGGUUGUAVAGU GA GG C           GG CCC UUC AUCCUCCGGCAUCA CU CC A           A CU G - - - - - AGAGGAA C           </pre>

Fig. 7 (cont.)

let-7f-1	UGAGGUAGUAGAUUGUUAUGUU	<p>AGU GAGGUAGUAGAUUGUUAUGUU UG          UCAG GAGGUAGUAGAUUGUUAUGUU GGGUAG          AGUC UUCCGUUAUCUAACAUAUCAUA UCCCAU A          CC- GAGGACUUG UUU</p>
let-7f-2	UGAGGUAGUAGAUUGUUAUGUU	<p>U UCAU          CUGUGGA GAGGUAGUAGAUUGUUAUGUU UUAGG A          GGCACCCU UUCUGUCAUCUGACAUAUCA GGUUC C          UAGA ACCC</p>
let-7g	UGAGGUAGUAGUUUGUACAGUA	<p>A U UGAGG A- A A          CC GGC GAGGUAGU GUUUGUACAGUU GUCU UG UACC C          GG CCG UUCCGUCA CGGACAUGUCA UAGA AC AUGG C          A - C ----- GG - C</p>
let-7h	UGAGGUAGUAGUGUGUACAGUU	
let-7i	UGAGGUAGUAGUUUGUGCU	<p>U U UGUG          CUGGC GAGGUAGUAGUUUGUGCU GUU GG CGGU          GAUCG UUCCGUCAUCGACGCG CAA UC GCCCG A          U UAGAGGUG - UUAC</p>
miR-1	UGGAAUGUAAAGAGUUGGAG	<p>A UUUUGAGA C A - AUA          UUC GCC GUUCCAUUGUUC UUGCAUUC AUA GUU          GAG CGG CGAGGUAGUAG AUUGUAAG UAU CGA U          - UCUAAG A G A ACU</p>
miR-1b	UGGAAUGUAAAGAGUUGUAA	<p>A GC ---- AC          UGGGA ACAUACUUCUUUAUU CCAUA UGG          ACUCU UGUUUGAAGAAUGUA GGUAU AUC C          A- A- CGA GU          AL49263.5</p>

Fig. 7 (cont.)

miR-1c	UGGAUGUAAAGAAGUAGUAC	
miR-1d	UGGAUGUAAAGAAGUAGUAAU	<p>C GC UGAACC U</p> <p>GCUUGGGA ACAUACUUCUUUAU CCAUA U</p> <p>CGGACUUU UGUUUGAAGAAUGUA GGUAA G</p> <p>A- A- CGAAUC</p>
miR-2a-1	UAUCACAGCCAGCUUUGAUGGC	<p>A AUUC UU</p> <p>GCUGGGCUC UCAAAG UGGUUGUGA AUGC CGC \</p> <p>CGAUUCGAG AGUUUC ACCGACACU UAGG GCG U</p> <p>U G A ----- CG</p>
miR-2a-2	UAUCACAGCCAGCUUUGAUGGC	<p>A C GAUAC</p> <p>AUCU AGC UCAUCAAG UGGUUGUGAUAG \</p> <p>UAGG UCG AGUAGUUU ACCGACACUAUAC C</p> <p>A - CG GCAAC</p>
miR-2b-1	UAUCACAGCCAGCUUUGAUGGC	<p>U UG - A C---- U</p> <p>CU CAAC UCUUCAAG UGGC GUGA AUGUUG C</p> <p>GG GUGG AGGAGUUUC ACCG CACU UUAUAC A</p> <p>C CG G A AUACU A</p>
miR-2b-2	UAUCACAGCCAGCUUUGAUGGC	<p>A A UUU--- CUU</p> <p>UUGUGUC UUCUCAAAG UGGUUGUGA AUG GC U</p> <p>AGCGCAG GAGGAGUUUC ACCGACACU UAC CG U</p> <p>C G A UUAUC UAU</p>
miR-3	UCACUGGGCAAAGUGUGUCUCA	<p>C C G U UUCA</p> <p>GAUC UGGGAUGCAU UUGU CAGU AUGU \</p> <p>CUAG ACUCUGUGUG AACG GUCA UACA A</p> <p>A A G C CUCU</p>

Fig. 7 (cont.)

miR-4	AUAAGCUAGACACCAUUGA	U UU C C C GG UU UUGCAU AGUUC UGU GUC AGC UUA UGAUU \ GGUGUUG UUGAAG ACCA CAG UCG AAU ACUGG U C UU A A A -- CC
miR-5	AAAGGAACGAUCGUUGUGAUUG	UA--- C AGUUGU \ GC AAAGGAA GAUCGUUGUGAUUG \ CG UUUCCUU UUAGUGACACUAUAC U CAAUA - AAUCCU
miR-6-1	UAUCACAGUGGCUGUUCUUUUU	A- C AG UAAUA \ UUUA UGUAGAGGGAUAGUUGCUGUG UGUA U \ AAAU AUGUUUUUCUUGUCGGUGACAC AAUA A U CC U CU UACCA
miR-6-2	UAUCACAGUGGCUGUUCUUUUU	C UU UG C U - G UAACC AAGGAAAC C CUG UGAUUA UA UU A GUUGG UUUUCUUG G GAC ACUAUAU AU AA A U UC GU C C A
miR-6-3	UAUCACAGUGGCUGUUCUUUUU	A A U AAAC \ CAA A GAAGGGAACGGUUGCUG UGAUGUAG UUG \ GUUU UUUUUUCUUGUCGGUGAC ACUAUAU AAC U G - U ACUC
miR-7	UGGAAGACUAGUGAUUUUGUUGU	U U U U -- UGGUC \ GAGUGCAU CCGUA GGAAGAC AG GAUUU UGUUGUU \ UUUACGUG GGCAU UCUUCUG UC CUAAA ACAUAAA U C - U C UA UGGUU
miR-8	UAAUACUGUCAGGUAAGAUGUC	CUGUUC - G C UCCUUU \ AAGGACAU ACAUCUU ACC GGCAG AUUAGA \ UUCCUGUG UGUAGAA UGG CUGUC UAAUCU U CCUGC- A A A CAAUAU

Fig. 7 (cont.)

miR-9	UCUUUGGUUAUCUAGCUGUAUGA	<p> <u>U</u> <u>UAU</u> <u>G</u> - GAU              GCUA UGUUG CUUUGU CUAGCU UAUGA GU A              CGAU AUAUU GRAGCCA GAUCGA AUACU CA A              U U UUC A G AUA         </p>
miR-10	ACCCUGUAGAUCCGAAUUUGU	<p>             CU - G U AUACU              CCACGU ACC CU UAGA CCGAAUUUGUUU A              GGUGUG UGG GA AUCU GCCUUAACACAGGA G              UU A G U AUUUC         </p>
miR-11	CAUCACAGUCUGAGUUCUUGC	<p>             U UCU CCC U ACU              GCACUUG CAAGRACUU CUGUGA GCG GU U              CGUGAGU GUUCUUGAG GACACU CGC CG A              C UCU A-- - AAA         </p>
miR-12	UGAGUAUUACAUCAGGUACUGGU	<p> <u>UG</u> <u>U</u> <u>C</u> - GCCUU              UACGGU AGU<u>U</u> AC<u>U</u> AGGUACUGGU GU A              GUGCCG UCA<u>U</u> UGU<u>U</u> UUCAUGACCA CA A              CA C - A ACCUA         </p>
miR-13a	UAUCACAGCCAUUUUGAUGAGU	<p>             U C - A UC-- CU              UACG AACUC UCA<u>U</u> AGGUGUGA AUG GA A              GUGC UUGAG AGUUUU CCGACACU UAC CU U              U U A A A UCAU AU         </p>
miR-13b-1	UAUCACAGCCAUUUUGACGAGU	<p>             UG- U ACU UAUU              CCA UCGUUA<u>U</u>AAUG UUGUGA UAUG C              GGU AGCAGUUU<u>U</u>AC GACACU AUAC A              UUG C --- UAAC         </p>
miR-13b-2	UAUCACAGCCAUUUUGACGAGU	<p>             UAUU G A GCUA UU              AAC CGUCA<u>U</u>AAUG CUGUGA UGUGGA U              UUG GCAGUUU<u>U</u>AC GACACU AUACUU G              GU-- A C ---- CA         </p>

Fig. 7 (cont.)

miR-14	UCAGUCUUUUUCUCUCUCCUA	<p>C C C C GCUU                  UGUGGGAG GAGA GGGGACU ACUGU                  AUAUCCUC CUCU UUUUCUGA UGAUA A                  U U C AAUU</p>
miR-15a	UAGCAGCACAUAAUUGGUUUGUG	<p>GAGUAAAGUA UA GA U                  CCUUG GCAGCACA AUGGUUUGUG UUU                  GGAAC CGUCGUGU UACCGGACGU AAA G                  AUA AAAACUC UA GG A</p>
miR-15b	UAGCAGCACAUCAUGGUUUACA	<p>U C C A A ACA                  CUG AGCAGCA AU AUGGUU CAU CU                  GAU UCGUCGU UA UACUAAG GUA GA G                  C U U C - ACU</p>
miR-16	UAGCAGCACGUAAAUUUGGCG	<p>AG C - A CGUUA UCUA                  GUCAGC UGC UUAGCAGCAC GU AAUAUUGG AGAU                  CAGUUG AUG AGUCGUCGUG CA UUAUGACC UCUA A                  GA A U A ----- UUA</p>
miR-16	only different precursor	<p>UC CU UA C AG AAU                  GU CACU AGCAGCAGG AAUAUUGG GU UGA A                  CA GUGA UCGUCGUGU UUAUAUACC CA AUU U                  GU UU CA A A- AUA</p>
miR-17	ACUGCAGUGAAGGCACUUGU	<p>GA CA- A G G - AUA                  GUCA AUAUUGU AAGUCUU CA UGCAG UAG UG                  CAGU UAUUACG UUCACGGGA GU ACGUC AUC AC U                  GG AUG A G - U GUG</p>
miR-18	UAAGGUGCAUCUAGUGCAGAUUA	<p>CU U C U A UGAA AG                  UGUU AAGG GCAU UAG GCAG UAG GU A                  ACGG UUCC CGUG AUC CGUC AUC CG U                  UC U A C - UA-- AU</p>

Fig. 7 (cont.)

miR-19a	UGUGCAA <u>AAUCUAUGCAA</u> AAACUGA	U U GCAG CC CUGUAGUUUGCAUAG UUGCAC UACA \ AGA CGUC GG GGUAGUCAAAACGU <u>AUC AACGUG</u> AUGU A C U UA UUG AAG
miR-19b-1	UGUGCAA <u>AAUCCAUGCAA</u> AAACUGA	UU UC UGUGUG CACUG CUAUGGUUAGUUUGCA GG UUUGCA CAGC \ GUGAU GGUGUCAGUCAAAACGU CC AAACGU GUCG A -- A U -- UCUUAU
miR-19b-2	UGUGCAA <u>AAUCCAUGCAA</u> AAACUGA	CUAC U ACAUUG U <u>UACAAU</u> AGUUUGCA GG UUUGCAU GCGUAUA A UGUA <u>AAU AGUU</u> AGUCAAAACGU CC AAACGUG UGUUAU U ---- A U UCGG G
miR-20	UAAAGUCUU <u>AAUAGUCAGG</u> UAG	C A- G - UU GUAG ACU AAGUCUU <u>AAUAGUCAG</u> UAG UG U CGUC UGA UUCACGAGUA <u>UUACGUC</u> AUC AU A A AA - U UG
miR-21	UAGCUUAUCAGACUG <u>AAUUGA</u>	A A A U AA UGUCGGUAGCUUAUC GACUG UGUUG CUGU G \ ACAGUCUGUCGGGUG CUGAC ACAAC GGUA C U - C - UC
miR-22	AAGCUGCCAGUUUGAAGAACUGU	U CC A U CCUG GGC GAG GCAGUAGUUUCUUCAG UGGCA GCUUUA GU \ CCG CUC CGUUGUCAAGAGUU ACCGU CGAAAU CG A U C- G - ACCC
miR-23a	AUCACA <u>UUGCCAGGAAU</u> UCC	C C G G CUUC GG CGG UGGGG U <u>UCCUGG</u> GAUG GAUUUG C CC GCC ACCUU AGGGACC UUAC CUA <u>AAAC</u> U A A U G A ACUG

Fig. 7 (cont.)

miR-23b	AUCACAUUGCCAGGGGAUUACCAC	<p>C U -- -- C GUGACU  GG UGC UGG GUUCCUGGCA UG UGAUUU U  CC ACG ACC UAGGGACCGU AC ACUAAA G  A C AU U -- AUUAGA</p>
miR-24-1	UGGCUCAGUUCAGCAGGAACAG	<p>G G A UA UCUCAU  CUCC GU CCU CUGAGCUGA UCAGU \ U  GAGG CA GGA GACUUGACU GGUCA U  A A C C- C- CACAUU</p>
miR-24-2	UGGCUCAGUUCAGCAGGAACAG	<p>CC CG CU- AA-- UU  CUCUG UCC UGC ACUGAGCUG ACACAG \  GGGAC AGG ACG UGACUCGGU UGUGUU G  A- -- ACU CACA UG</p>
miR-25	CAUUGCACUUUGUCGGUCUGA	<p>A AG G UU G UG ACG  GGCC GUGUUG AGGC GAGAC G GCAAU CUGG C  CCGG CGUGAC UCUG CUCUG C CGUUA GGUC U  C AG G UU A CG CCG</p>
miR-26a	UUCAAGUAUUCAGGAUAGGCU	<p>- G U U GCAG  AGCC GUG CCUCGU CAAGUAA CCAGGAUAGGCUGU G  UCCGG CGC GGGCA GUUCAUU GGUUCUAUCCGGUA U  G A C - ACCC</p>
miR-26b	UUCAAGUAUUCAGGAUAGGCU	<p>GA - U UC UGUG  CCGG CCC AGU CAAGUAAU AGGAUAGGUUG \  GGCC GGG UCG GUUCAUUA UCUUGUCCGAC C  AG C - CC CUGU</p>
miR-27a	UUCACAGUGGCCUAAGUUCGGCU	<p>A A A U G UCCAC  CUG GG GC GGGUUGAGCUGU GUGAGCA GG \  GAC CC CG CUUGAAUCCGGUGA CACUUGU CU A  C C C - G GAACC</p>



Fig. 7 (cont.)

miR-27b	UUCACAGUGGCCUAAGUUCUG	<p>AUUG UGAU U            AGGUCAGAGCUUAGCUG GUGAACAG UGG \</p> <p>UCCACGUCUUGAAUCGGU CACUUGUU GCC U</p> <p>GA-- UC-- U</p>
miR-28	AAGGAGCUCACAGUCUAUUGAG	<p>C A U ----- CC</p> <p>GGU CUUGCCUC AGGAGCUCACAGUCUA UG AGUUA U</p> <p>UCA GGACGGGAG UCCUGGAGUUGUAGU AC UCAGU U</p> <p>C G C CCUU CU</p>
miR-29a	CUAGCACCAUCUGAAAUCGGUU	<p>UUU C UCAAU</p> <p>AUGACUGAUUUC UGGUUU AGAG \</p> <p>UAUUGGCUAAAG ACCACGA UCUU A</p> <p>UCU - UUAUU</p>
miR-29b	UAGCACCAUUUGAAAUCAGUGUU	<p>A U GU UUAUU</p> <p>AGGA GCUGGUUCA AUGGUG UUAGAU \</p> <p>UCUU UGACUAAAGU UACCAC GAUCUG A</p> <p>G U -- -- UUAGUG</p>
miR-29c	UAGCACCAUUUGAAAUCGGUua	
miR-30a-s	UGUAAAACAUCUCGACUGGAAGC	<p>A UC ----- A</p> <p>GCG CUGUAAAACAUC GACUGGAAGCU GUG A</p> <p>CGU GACGUUUUGUAGG CUGACUUUCGG CAC G</p> <p>C -- -- GUAAA C</p>
miR-30a-as	CUUUCAGUCGGGAUGUUUGCAGC	<p>A UC ----- A</p> <p>GCG CUGUAAAACAUC GACUGGAAGCU GUG A</p> <p>CGU GACGUUUUGUAGG CUGACUUUCGG CAC G</p> <p>C -- -- GUAGA C</p>

Fig. 7 (cont.)

miR-30b	UGUAAACAUCUACACUCAGC	<p>U - UCAUA  <u>AUGUAAACAUC</u> <u>ACA</u> <u>CUCAGCUG</u> C          UGCAUUUGUAGG UGU GGGUCGGU A          - A UGCGU</p>
miR-30c	UGUAAACAUCUACACUCUCAGC	<p>UACU U ACA GUGGAA          AGA <u>GUAACA</u> <u>CCU</u> <u>CUCUCAGCU</u> A          UCU CAUUUGU GGA GAGGGUCGA G          UUCU C A-- AAGAAU human</p>
miR-30d	UGUAAACAUCUACACUCUGAAG	<p>U U CCC GUAAGA          GU GU <u>GUAACAUC</u> <u>GACUGGAAGCU</u> C          CA CG CGUUUGUAG CUGACUUUCGA A          U U A-- AUCGAC chr8 human</p>
miR-31	GGCAAGAUCUGGCAUAGCUG	<p>GA G C U- GAA          GGAGAG <u>GGCAA</u> <u>AUG</u> <u>UGGCAUAGC</u> GUU C          CCUUUC CCGUU UAC ACCGUAUCG CAA U          UA A A UC GGG</p>
miR-32	UAUUGCACAUUACUAAGUUGC	<p>U - UU C          GGAGAUUUGCACAU <u>ACUAAGUUGCAU</u> G GU A          CUUUUAUAGUGUGUG UGAUUUAACGUA C CG C          - A UC G</p>
miR-33	GUGCAUUGUAGUUGCAUUG	<p>A UU UUCU UG          CUGUGGUGCAUUGU G <u>GCAUUGCAUG</u> GG          GACACUACGUGACA C UGUACGUAC CC G          C UU ----- AU</p>
miR-99a	ACCCGUAGAUCCGAUCUUGU	<p>A UC U G AAG          CAUA <u>ACCCGUAGA</u> <u>CGA</u> <u>CUUGUG</u> UG U          GUGU UGGGUAUCU GCU GAACGC GC G          C UU C - CAG</p>

Fig. 7 (cont.)

miR-99b	CACCCGUAGAACCGACCUUGCG	<p>CC AC C ----- C</p> <p>GGCAC ACCCGUAGA CGA CU UGCGG GG \</p> <p>CUGUG UGGGUGUCU GCU GA ACGCC CU C</p> <p>CC GU C ACAC G U</p>
miR-101	UACAGUACUGUGAUAACUGA	<p>A GUCCA</p> <p>UCAGUUAUCACAGUGCUG UGCU U</p> <p>AGUCAUAUGUGUCAUGAC AUGG U</p> <p>AAAUUC</p>
miR-122a	UGGAGUGUGACAUAUGGUGUUUGU	<p>GG C UGUCC</p> <p>AGCUGU AGUGUGA AAUGGUGUUUG A</p> <p>UCGAUA UCACACU UUAACCGCAAC A</p> <p>AA A UAUCA</p> <p>woodchuck</p>
miR-122b	UGGAGUGUGACAUAUGGUGUUUGA	
miR-122a, b	UGGAGUGUGACAUAUGGUGUUUG	
miR-123	CAUUAUUACUUUUGGUACGCG	<p>A A U CGCUG C</p> <p>UGAC GC CAUUAUUACUU UGGUACG UGA A</p> <p>ACUG CG GUAAUAUAUGAG GCCAUGC ACU C</p> <p>G C U UCAA- U</p>
miR-124a*	UUAAGGCACGCGGUGAAUGCCA	<p>- C A GA UAAUG</p> <p>CUCU G GUGUUCAC GCG CCUUGAUU U</p> <p>GAGA C CGUAAGUG CGC GGAUUUA C</p> <p>A - G AC CAUUAU</p>

Fig. 7 (cont.)

miR-124b	UUAAGGCACGGGGUGAAUGC	<p>CC A GA UAAUG          CUCU GUGUUCAC GCG CCUUGAUU \          GAGA CGUAAGUG CGC GGAUUAA U          AC G AC CAUAC AC021518</p>
miR-125a	UCCUGAGACCCUUUAACCUGUG potential lin-4 ortholog	<p>C C UA ---- A          CUGGGU CCUGAGA CCUU ACCUGUGA GG C          GGUCCG GGGUUCU GGAG UGGACACU CC G          A U -- GGGA U</p>
miR-125b	UCCUGAGACCCUAACUUGUGA potential lin-4 ortholog	<p>UC C A GG- U          GCCUAG CCUGAGA CCU ACUUGUGA UAU U          CGGAUC GGGUUCU GGA UGARACACU AUG U          CA U C ACA A</p>
miR-126	UCGUACCGUGAGUAAUAUGC	<p>A U CGCUG C          GC CAUUUUACUU UGGUACG UGA A          CG GUAAUAUAGAG GCCAUGC ACU C          C U UCAA- U</p>
miR-127	UCGGAUCCGUCUGAGCUUGGCU	<p>A U G G C -- AG          CC GCC GCU AAGCUCAGA GG UCUGAU UC \          GG UGG CGG UUCGAGUCU CC AGGCUA AG A          C U - - G U CU AA</p>
miR-128	UCACAGUGAACCGGUCUCUUUU	<p>UUC UAG CU U          GUUGGA GGGGCCG CACUGU GAGAGGU U          CGACUU CUCUGGC GUGACA CUCUUUA A          UUU CAA -- C</p>
miR-129	CUUUUUUCGGUCUGGGCUGC	<p>- C CU UCCU C          GGAU CUUUUUG GGU GGGCUU CUG CU A          UCUA GAAAAAC CCA CCCGAA GAC GA A          U C UU G UGAU- C human</p>

Fig. 7 (cont.)

miR-130	CAGUGCAAUUGUUAAAAGGGC	<p>- C A GUCUAAC GA GCUCUUU ACAUUGGCU CU \ G CU CGGAAA UGUAACGUGA GA G A U C GCCAUGU</p>
miR-131	UAAAGCUAGAUAAACCGAAGU	<p>G C G U A GUU UUAU UUUGGUUAUCUGCU UAUGAG GU U CAA AAUG AAGCCAAUAGAUGCA AUACUU UG U A A A C G</p>
miR-132	UAAAGUCUACAGCCCAUGGUGU	<p>A UUC G- G GGC ACCGUGGCU GAUUGUACU UGG \ G CCCG UGGUACCGA CUGACAAUGG GCC A C CAU AG A</p>
miR-133	UUGGUCCCCUUAACCAGCUGU	<p>A AA U A GCCUC GCUA AGCUGU AA GG ACCRAAUC U CGAU UCGACCA UU CC UGGUUUAG U G AC C C GUAAC</p>
miR-134	UGUGACUGGUUGACCAGAGGGA	<p>GU U A- G GCGU AC AGGU GUGACUGG UG CCA AGGG GC \ G UCCCA CACUGAUC AC GGU UCCE UG U AC C CG G ACU- UC</p>
miR-135	UAUGGCCUUUUUAUCCUAUGGAA	<p>UU UUCUUAU CUAUGGCCUUU AUUCCUAUGUGA \ U GGUGCCGAGG UAGGGAUAUACU U U- CGCUCG</p>
miR-136	ACUCCAUUUGUUUUGAUGAUGGA	<p>C UUU UUCU GAGGACUC AUUUG UGAUGAUGGA \ U CUUCUGAG UAAAC GCUACUACCU U - UCU CGAA</p>

Fig. 7 (cont.)

miR-137	UAUUGC <u>UUAAGAUAACGGUAG</u>	<p>G G A - GA          CUUCGGU ACG <u>GUUUCUUGGGUGG</u> UAAUA CG \</p> <p>GGAGCUG UGC <u>CAUAGAUAUCGUU</u> AUUGU GC U</p> <p>A G - - U AU</p>
miR-138	AGCUGGUGUUGUGA <u>UVC</u>	<p>-- UCA AC- C CG          CAGCU GGUGUUGUGAA GGCCG GAG AG C</p> <p>GUUGG CCACAGCACUU UCGGC UUC UC A</p> <p>GA UA- CCA - CU</p>
miR-139	UCUACAGUGCACGGUG <u>UCU</u>	<p>G - U A GUGGC          GU UAUCUA CAG GC CGUGUCUCCAGU \</p> <p>CA AUGAGGU GUC CG GCGCAGAGGUCG U</p> <p>- U C - GAGGC</p> <p>human</p>
miR-140	AGUGGUUUUACCCUAUGG <u>UAG</u>	<p>- A A A UU UC          CCUG CC GUGGUUUACCCU UGGUAGG ACG A</p> <p>GGAC GG CACCAAGAUGGGA ACCAUCU UGU U</p> <p>A - - C -- CG</p>
miR-141	AACACUGUCUGGUA <u>AAAGAUGG</u>	<p>U -- U AU GAAG          GGG CCAUCUU CCAG GCAGUGUUGG GGUU \</p> <p>CCC GGUAGAA GGUC UGUCAACAUC UCGA U</p> <p>- - AU - C- AGUA</p>
miR-142s	CAUAAAGUAGA <u>AAAGCACUAC</u>	<p>AC- A UAA--- G          CCAUAAAGUAG AAGCACUAC CA C</p> <p>GGUAUUUCAUC UUUGUGAUG GU A</p> <p>GUA C UGGGAG C</p>
miR-142as*	UGUAGUGUUUCCUACUUU <u>AUGG</u>	<p>AC- A UAA--- G          CCAUAAAGUAG AAGCACUAC CA C</p> <p>GGUAUUUCAUC UUUGUGAUG GU A</p> <p>GUA C UGGGAG C</p>

Fig. 7 (cont.)

new	AUAAGACGAGCAAAAAGCUUGU	<p>G C G G C AU</p> <p>UGAC GCGGAGCUUU GC CG UUAUAC UG \</p> <p>ACUG UUGUUCGAAA CG GC AAUAUG AC G</p> <p>G A A AG C UC</p> <p>AL049829.4</p>
miR-143	UGAGAUGAAGCACUGUAGCuca UUAGAUGAAGCACUGUAG	<p>G G U - AG</p> <p>CCUGAG UGCAGUGCU CAUCUC GG UC U</p> <p>GGACUC AUGUCACGA GUAGAG CU AG U</p> <p>G A U G GG</p> <p>AC008681.7</p>
miR-144	UACAGUAUAGAUGAUGACUAG	<p>G A A- GU</p> <p>GGCUGG AUAUCAUC UUAUCUGUA GUUU G</p> <p>CUGAUC UGUAGUAG AUAUGACAU CAGA A</p> <p>A - CA GU</p>
miR-145	GUCCAGUUUCCCGAGAAUCCCUU	<p>C UC U C UGGAUG</p> <p>CUCA GG CAGU UU CCAGGAUCCCU \</p> <p>GAGU UC GUCA AA GGUCCUAGGGG C</p> <p>- UU U A UAGAAU</p>
miR-146	UGAGAACUGAAUCCAUUGGGUUU	<p>CU C AUAUC</p> <p>AGCU GAGAACUGAAUU CAUGGGUU A</p> <p>UCGA UUCUUGACUUA GUGUCCAG A</p> <p>C- A ACUGU</p>
miR-147	GUGUGUGGAAAUUGCUCUGCC	<p>A- CAA ACA--- GA</p> <p>AAUCUA AGA CAUUUCUGCACAC CCA \</p> <p>UUAGAU UCU GUAAAGGUGUGUG GGU C</p> <p>CG UC- ACCGAA AU human</p>
miR-148	UCAGUGCACUACAGAACUUUGU	<p>- A- CC - AGU</p> <p>GAGGCAAAGUUCUG AG CACU GACU CUG \</p> <p>CUCUUUUCAAGAC UC GUGA CUGA GAU A</p> <p>A AC -- A AGU human</p>

Fig. 7 (cont.)

miR-149	UCUGGCCUCCGUGUCUUCACUCC	<p style="text-align: center;"> <u>G C G A</u>    <u>GUG G</u>                  GGCUCUG CUC GU UCUUC CUCCC UUU U                  UCGGGC GAG CA GGAGG GAGGG GAG C                  G A G - AG- C             </p>
miR-150	UCUCCCAACCCUUGUACCAGUGU	<p style="text-align: center;"> <u>AC U</u>    <u>UG- UG</u>                  CCCUGUCUCCA CCU GUACCAG CUG \                  GGGUAVAGGGGU GGA CAUGGUC GAC C                  CC - CCA UC             </p>
miR-151	CUAGACUGAGGCCUCCUUGAGGU	<p style="text-align: center;">                 C CA UGUCU                  CCUG CCUCGAGAGCU CAGUCUAGUA \                  GGAC GGAGUCCUCGG GUCAGAUCAU C                  A A- CCCUC             </p>
miR-152	UCAGUGCRAUGACAGAACUUGG	<p style="text-align: center;">                 G A CC CGG C                  CCGGGCCUAGGUUCUGU AU CACU GACU GCU U                  GGCCCGGGUUCAGACA UA GUGA CUGA CGA G                  G C - - - - G             </p>
miR-153	UUGCAUAGUCACAAAAGUGA	<p style="text-align: center;">                 GU A- AAU                  CAGUG UCAUUUUUGUGAU UGCAGCU GU \                  GUUAC AGUGAAAACACUG ACGUUGA CG A                  U AU CC AGU             </p>
miR-154	UAGGUUAUCCGUGUUGCCUUCG	<p style="text-align: center;">                 U - CCU-- UUU                  GAAGAUAGGUUA CCGUGU UG UCGC \                  UUUUUUACCAGU GGCACA AC AGUG A                  U U UAAGC UUU             </p>
miR-155 [BIC-RNA]	UUUAUGCUAAUUGUGAUAGGGG	<p style="text-align: center;">                 U U A UUGGCC                  CUGUUAUUGCUAAU G G UAGGGGU \                  GACAAUACGAUUG U C AUCCUCAG U                  - C - UCAGUC             </p>



Fig. 7 (cont.)

name	sequence	structure
miR-C1	AACAUUCAACGCUGUCGGUGAGU	<pre> U A U CU A GGGAUUCA CCA GG ACA UCAACG GUCGGUG GUU GGU CC UGU AGUUGC CAGCCAC CAA A U A C -- - AAAACAAA                     </pre>
miR-C2	UUUGGCAUUGGAGAACUCACA	<pre> UU UGG UCA UAAGGU ACCAU UUGGCAA UAGAAC CACCGG A UGGUA AACCGUU AUCUUG GUGGCC A UC CAG --- CAGGGU                     </pre>
miR-C3	UAUGGCACUGGUAGAUAUCACUG	<pre> G AC-- GA -- AC CUGU UAUGGC UGGUA AUUCACUG UGA A GACA AUACCG GCCAU UAAGUGAC ACU G A GGAA -- UG CU                     </pre>
miR-C4	CUUUUUGCGGUCUGGGCUTUGUU	<pre> -- C CU G UUUU C UGGAU CUUUUUG GGU GGGUU CUG CU G AUCUA GAAAAAC CCA CCCGAA GAC GA A U C C UU G UGAU C                     </pre>
miR-C5	UGGACGGAGAACUGAUAAGGGU	<pre> U C C AG - UG CCU UCCUUAUCA UUUUCC CCAGC UUUG A GGA GGGAAUAGU AAGAGG GGUUG GAAU C U C CA U CU                     </pre>
miR-C6	UGGAGAGAAAGGCAGUUC	<pre> A G AU UC AGGAUUGGAG GAAAG CAGUUCUUG GG C UUCCUGGUCUC CUUUC GUCGGGGAC CC C - G -- UC                     </pre>

Fig 7 (cont.)

name	sequence	structure
miR-C7	CAAAGAAUUCUCCUUUUGGGCUU	<pre>           U  UU  UCUCAU ACUUUCCAAAGAAUUC CCUU GGGCUU  U UGAAGGGUUUUUUUAAAG GGAA CCCGAA  U           U  U-  UUUUUAU           </pre>
miR-C8	UCGUGUCUUGUGUUGCAGCCGG	<pre>           A  A  C  CGCUGC UC GGCU CAACACAGGAC CGGG  U GG CCGA GUUGUUCUG GCUC  C           C  U  CCCAGU           </pre>
miR-C9	UAAACAGUCUGGUAACGAUGU	<pre>           C  UU  UUG GGGCAUC UUAACCGACAGUG UGGA UC \ CUUGUAG AAUGGUCUGUCAGAUUCU AG G           C  A  C-  UUC           </pre>
miR-C10	CAUCCCUUGCAUGGUGGAGGGU	<pre>           CA UC  GU  UGAGCUC UCU CA CCUUGCAUG GGAGGG  U AGG GU GGGACGUAC CCUCCC  C           AC UU  AC  CAAAAGU           </pre>
miR-C11	GUGCCUACUGAGCUGACAUCAGU	<pre>           G  G  A  UA  UCUCAU CUCC GU CCU CUGAGCUGA UCAGU \ GAGG CA GGA GACUUGACU GGUCA  U           A  A  C  C-  CACACU           </pre>
miR-C12	UGAUAUGUUUGAUAUAUUAGGU	<pre>           U-  UA---  UU CUGUG GAUAUGUUUGAUAUAU GGUUG \ GACAU UUAUACGAACUAUAUA CUAUA A           CC  UCAAC  UU           </pre>

Fig. 7 (cont.)

name	sequence	structure
miR-C13	CAACGGAAUCCCCAAAGCAGCU	<pre> C   C   AA   UU   -   C AGCGG AACGGAUCC AA GCAGCUG GU CU C UGGUCC UUGC UUAGG UU CGUCGAC UA GA A C   -   CA   CU   C   G                     </pre>
miR-C14	CUGACCUAUGAAUUGACA	<pre> C   -   A   A   UGCUCUC UGACCUAUG AAUUG CAGCCAG G ACUGGAUAC UUAAC GUCGGUC U -   C   C   UCCCCUC                     </pre>
miR-C15	UACCACAGGGUAGAACCCACGGA	<pre> -   G   A   A   UU   UC UCCUG CCG UGGUUUAACCCU UGGUAGG ACG A AGGAC GGC ACCAAGAUGGGA ACCAUCU UGU U A   -   -   C   -   -   CG                     </pre>
miR-C16	AACUGGCCUACAAAGUCCCGAG	<pre> A   U   C   A   A   AGU GAG GCUGGG CUUUG GGGC AG UGAG G CUC UGACCC GAAAC UCCG UC ACUU U C   U   A   G   A   GAC                     </pre>
miR-C17	UGUAAACAGCAACUCCAUUGUGGA	<pre> U   A   G   -   -   U AUCGGG GUAACAGCA CUCCAU UGGA CUG G UAGUCU CAUUGUCGU GAGGUG ACCU GGC C U   C   -   -   UA   U                     </pre>
miR-C18	UAGCAGCACAGAAAUUUGGC	<pre> U   A   -   UG   GAA AGCAGCACAG AAUADUGGCA GG G UCGUCGUGUC UUAUAACCGU CU U GG   -   -   GAG                     </pre>

Fig 7 (cont.)

name	sequence	structure
miR-C19	UAGGUGUUCUUGUUGUUGG	<pre> A  A  C  GGCUGGG GUGAAU  GGU  GDU  AUGUUGUUG  U CACUUG  CCA  CAA  UACAACAAC  U C  C  C  U  ACAAGUCU           </pre>
miR-C20	UUCACCACCUUCUCCACCAGC	<pre> C  A  CA  GA  -  A GGCUGGC  GGU  GAGAGG  GUGG  GGU  AAG  G CCGGUACG  CCA  CUCUUC  CACU  CCA  UUC  C A  C  C  AC  UC  C  U           </pre>
miR-C21	GGUCCAGAGGGAGAUAGG	<pre> G  -  C  G  U  UGCCUG UCAU  G  UC  A  AGGGAGA  AGG  U AGUAA  U  AG  U  UCUCUUCU  UCC  G A  A  A  A  -  UUUUUA           </pre>
miR-C22	CCCAGUUCAGACUACCUGUU	<pre> AAC  U  C  U  G---  G GCC  CCAGUGU  CAGACUAC  UGU  CA  GAG  \ CGG  GGUUACA  GUCUGAUG  ACA  GU  CUC  C AUU  C  -  U  GUAA  U           </pre>
miR-C23	UAAUACUGCCUGGUAUGAUGAC	<pre> GGC  -  C  UAGUG GCCGU  CAUC  UUACUGGGCAG  AUUGGA  U CGGCA  GUAG  AUUGGUCCGUC  UAAUCU  C ---  U  A  CUAGU           </pre>
miR-C24	UACUCAGUAAGGCAUUGUUCU	<pre> U  U  U  UUC  A UACCUAC  CAG  AAGGCAUUGUUC  UAU  U AUGGGAUG  GUC  UUCCGUGACAAG  AUA  U U  U  U  UAA  A           </pre>

Fig. 7 (cont.)

name	sequence	structure
miR-C25	AGAGGUAUAGCGCAUGGGAAGA	<pre> U      A-      UG      C GUUCC UUUUCCUAUGC UAUACUUCUU UGGAU \ CGAGG <u>AGAAGGGUJACG</u> <u>AUAUGGAGAA</u> AUCUG U       U      CG      --      G           </pre>
miR-C26	UGAAAUGUUUAGGACCACUAG	<pre> C      U      G      A      C      U GGUC AGUGGUUCU GACA UUCA CAGUU UG \ CCAG <u>UCACCAGGA</u> <u>UUGU AAGU</u> GUUAA AC A       A      U      A      -      C      G           </pre>
miR-C27	UUCUUUUUUGUCAUCCUAUGCCUG	<pre> U      A      U      GAGAUA UGGAC <u>UCCUUUUGUC</u> <u>UCCUA GCCU</u> \ ACUUG AGGAAAACGG AGGGU CGGA      U       C      A      GGAAGUA           </pre>
miR-C28	UCCUUCAUCCACCGGAGUCUG	<pre> UC      C      UCUUA CUCUUG <u>CUUCAUCCAC</u> <u>GGAGUCUG</u>      U GAGGAC <u>GAAGUGAGGUG</u> <u>CUUAGAC</u>      G       UC      A      CAACC           </pre>
miR-C29	GUGAAUGUUUAGGACCACUAGA	<pre> U      C      U      G      A      C      U GCC GGUC AGUGGUUCU GACA UUCA CAGUU UG \ CGG <u>CCAG UCACCAGGA</u> <u>UUGU AAGU</u> GUUAA AC A       C      A      U      A      -      C      G           </pre>
miR-C30	UGGAAUGUAAGGAAGUGUGUGG	<pre>       C      U      AUAUC CCAGG CCACAUGCUCUUUAU C CADAG \ GGUUU <u>GGUGUGUAGGAAGUA</u> <u>G GUAUC</u>      U       U      A      -      ACCGAC           </pre>

Fig 7 (cont.)

name	sequence	structure
miR-C31	UACAGUAGUCUGCACAUUGGUU	<pre> AUC   U   C   -----   G GCC  CCAGUGU CAGACUAC UGU      UCAG A CGG  GGUUACA GUCUGAUG ACA      GGUC G AUU   C   -   UGUACAG   G           </pre>
miR-C32	CCUGUAGAACCGAAUUUGUGU a miR-10 variant	<pre> A   G   C   UG- AC UAUAU CCCU UAGAA CGAAUUUGUG GU C AUADA GGGG AUUU GCUUAGACAC UA C A   -   A   UGA CA           </pre>
miR-C33	AACCCGUAGAUCCGACUUGUGA A a miR-99a variant	<pre> A   C   C   A   C   AU CACA ACC GUAGAU CGA CUUGUG UG U GUGU UGG UAUCUG GUU GAACAC AC C A   A   U   C   -   GU           </pre>
miR-C34	GCUUCUCCUGGCUCUCCUCCUCCUC	<pre> C   U   UUG   GGAG AAGG AGGGG GAGGG CGGGAGGAGC CGGGC G UUCC UCUCU CUCCUC GUCCUCUCCG GUUCG C -   -   UCG   C   CCGU           </pre>



Fig. 7 (cont.)

let-7l	miR-1	miR-1b	miR-1c	miR-1d	miR-2a-1	miR-2a-2	miR-2b-1	miR-2b-2	miR-3	miR-4
precursor ident. to mouse (AL117383.19); also AC048341.22		U97405.1 nt 1-22 (216)	U97405.1 nt no mouse hit (only nti-21)	AL48263.5 chr20 nti-22 (216)						
found, supported by EST BB661269	found	found	found	found						
				found						



Fig. 7 (cont.)

miR-5																				2R, AE003795
miR-6-1																				2R, AE003795
miR-6-2																				2R, AE00379
miR-6-3																				2R, AE00379
miR-7	AC003791 chr19 diff. precursor; EST BF371391 again different																			2R, AE003791
miR-8																				2R, AE003805
miR-9	AC005316 chr15 AC026701, chr3 each with diff. precursor									AP155142.1 chr19 diff prec, align diff prec.s in trace hits										3L, AE003516
miR-10	AF287967 chr11 (HOX B4/B5)																			AE001574
miR-11																				3R, AE003735
miR-12																				X, AE003499
miR-13a																				3R, AE003708

not cloned, but mouse EST predicts precursor similar to human

not found, but AC011194 chr.11 predicts diff. precursor

2diff  
precurs  
scaffold  
3868 and  
2417

found

Fig. 9 (cont.)

miR-13b-1											3R, AE003708			
miR-13b-2											X, AE003446			
miR-14											2R, AE003833			
miR-15a			13, AC069475				found	trace#72 137197 prec silyg diff						
miR-15b								trace#79 105069						
miR-16			13, AC069475 interesting leukemia locus				found						AL606727 diff precurs	
miR-16			3, NT_005710.6			several trace, near ly ident precursor			found trace#7910506 similarly ident prec. as in human					
miR-17			13, AL138714											
miR-18			13, AL138714											
miR-19a			13, AL138714											
miR-19b-1			13, AL138714						found				646757 with a U9C	



Fig. 7 (cont.)

miR-27a	19, AC020916	found	found, but no db hit	found, but no db hit for mouse	found	found	found	found				
miR-27b	XW_098943.1 chr9 identical precursor						found, maps to chr 13 NGSC mmtrace 44671617					
miR-28	3, AC063932											
miR-29a	7, AF017104 second ident. copy found in chr7 CLUSTER, this cluster also consvd in mouse; AC024913.32	found, AC024913.32	found, mmtrace#23467334	nearly ident precursor trace#23467334, EST AC024913.32			trace, EST, nearly ident prec	found				
miR-29b	AL035209.1 chr1 CLUSTER of miR-29-b and 29-c; miRNA similar to miR-83	found		AC024913.32, d iff precursor in EST, BG342396 (retina)			FOUND				Scaffold_17670. (A third copy)	
miR-29c				found			found, supported by ESTs	found			Scaffold_17670 has two copies of this RNA	
miR-30a-s	nearly ident fold in AL035467.23 chr6			found			found	found				
miR-30a-as	6, AL035467											
miR-30b	human AF159227.6 chr8, different precursor			found with diff. precursor in trace #85261735							Scaffold_3483, diff precursor	
miR-30c	AL136164.8 Chr. 6 supported by ESTs (BF594736.1)			found, but no db hit for mouse				found				

Fig. 7 (cont.)

mir-30d	AF159227.5 chr8						found, but no mouse ab hit	Scaffold 3483, diff fold
mir-31	9, AL353732							
mir-32	9, AL354797							
mir-33	22, 299716							G44780 with diff. precursor
mir-99a	AF000962.2 chr21, ident to mouse; (similar to miR-10 and miR-51)		trace#4891071					
mir-99b	AC018755.3 chr.19; (similar to miR-10 and miR-51)		mmtraos #92340982					
mir-101	AL158147.17 chr9 diff precursor		AK021368.1 cDNA eyeball	found				US3213.1 T.fluviat 1115
mir-122a					abundant but no db hit, except woodchuck X13234			
mir-122b								
mir-122a,b								
mir-123					genomic hits (trace#6108 147), no EST			Scaffold_3295

Fig. 7 (cont.)

miR-124a*	nearly ident. precursor in chr8(AC021518) chr20(AL096828) chr20(AL096828)	found in chr1V intron,diff precursor	found	found	most abundant in cereb.,genomic hits (tracef21097008, hits;precurs=11737241). found, but no db hit	abundant;seve ral.trace corebellum	found	found	slightly diff precursor AC009251 chr2L		
miR-124b	AC021518 chr8,nearly ident chr20 AL096828.29										
miR-125a	ident precu in AC018755.3 chr 19				genomic hits tracef33921945, 48262259 and more	found					
miR-125b	AP001359.4 chr11 AP001667.1 chr2L(chr2L like mouse)				mmtracef3521597 and more	tracef8398570 5	found with A22U	found in AC006590.1 1 with diff fold	Scaffold_2358		
miR-126											
miR-127	human AL117190.6 chr.14 same precurs as in mouse				hit in tracef79514537						
miR-128	ident in AC016742.10 chr. 2;diff prec in AC016943.7 chr.3				genomic hit tracef51670230	found	found		Scaffold_828,diff_prec		
miR-129	human AC018662.3 chr7				found, but no db hit						
miR-130					mmtraceq 68479278						with diff fold AC091299.2
miR-131	AC005317.2 chr 15 sligh.diff precursor,but AC026701.6 chr 5 ident				several trace hits,mouse AP155142	found					
miR-132	AL137038.5 chr17 prec sligh.diff from mouse				trace hit#86984641						

Fig. 7 (cont.)

mir-133	AL131221.15 chr6 diff. Precursor(ident to rat L33722.1)				found, trace# 62407955	found	AC093440.1 diff. Precursor	Scaffold_1049;prec u nearly like mouse	
mir-134	AL132709.5 chr14 similar precursor				trace#6462031 1				
mir-135	AC092045.2 chr3 AC018659.35 chr12 (ident or similar to mouse)				trace#7149523 5, EST#8780995 .1(kidn., sple en)(-chr3huma n)	found		Scaffold 2125 with similar precurs	
mir-136	AL117190.6 chr14 ident to mouse				trace#8607175 3				
mir-137	AC027691.1 chr1 , ident to mouse, nearly ident fish				trace#8977454 3, EST (hypothen.)#18 52436.1, ident			Scaffold_ 18244 nearly ident to mouse/man	
mir-138	AC006058.1 chr3 precursor diff				mouse EST 88528620.2				
mir-139	AP003065.2 chr11				found, but no mouse hit				
mir-140	AC026468.8 chr.16, precurs r nearly ident,			several trace hits; trace#1053 0393					
mir-141	AC006512.12 chr12, precursor slightli diff			AC002397 chr6		found			
mir-1425	AC004687.1 chr17 BCLJ/myq translocation locus, like mouse			found					
mir-142as*				several EST AY153235		found			

Fig. 7 (cont.)

new	AL049829.4 chr14								found but no db hit					
miR-143	AC008681.7 chr5			found, but no db hit	found				found					
miR-144	XM_064366.1 precursor nearly ident		found						EST AA990206 .1, trace 2143909					
miR-145	AC008681.7 chr5 GG->GA/precu nearly like mouse, see 2 positions above								found EST BF163348 .1 lung	Scaffold 934 similar				
miR-146	AC008388.7 chr5 diff precursor								trace#34 639321					
miR-147	AL592549.7								found					
miR-148	AC010719.4													
miR-149										found, no db hit				
miR-150														
miR-151														
miR-152	human chr 17 AC004777.1, nearly identical								trace#85 955550					
							trace#8472 1065,10352 801							
							trace#8845 6669							
													found in colon, supported by trace#8700445; close match MSC in chr18 (additional 14C unlikely, not supported by traces and	



Fig. 7 (cont.)

miR-153	AC006372.2 chr7 ident-precursor								found sever. mmtrace 87010874						
miR-154	AL132709.5 chr14 nearly identical precursor								found sever. mmtrace 86715639						
miR-155 {BIC-RNA}	human BIC RNA:AF402776.1 {has U12C}							found chr 16 mouse							

Fig. 7 (cont.)

name	human	mouse						Drosophila	fugu fish	zebratfish
		spleen	eye	kidney	testes	lung	thymus			
mir-C1	with different precursors in chr9 AL158075.11, chr1 AL136321.5		mouse trace #76647842						scaffold_1819	
mir-C2	chr7 AC084864.2 similar precursor		mouse trace #88841093						scaffold_967	AL590150.2
mir-C3	chr7 AC084864.2 ident-precursor		trace #86029980						scaffold_967	AL590150.2
mir-C4	similar precurs.in chr7 AC018662.3		trace #13885686		found					
mir-C5	chr15 AC069082.9		trace #87318220					found	scaffold_3671	
mir-C6	chr22 AC005664.2 ident-precursor		chr16 AC01326.12							
mir-C7	chr1 AL512443.7 similar prec.		trace #86694995							
mir-C8					found, trace #51673384					
mir-C9					found, trace #78964803				scaffold_2210, diff. precursor	
mir-C10	chrX AF222686.1 nearly ident. precursor				found, trace #61928192					
mir-C11	chr9 XM_098943.1 has C170; prec.nearly identical to mouse				found, cdNA AL286629.1, has C170					
mir-C12					found, trace#71760450				scaffold_2294	
mir-C13				found						





## SEQUENCE LISTING

<110> Max-Planck-Gesellschaft zur Förderung der Wissensc

<120> Identification of novel genes encoding for small  
temporal RNA

<130> 26250PWO\_DR

<140>

<141>

<150> EP 01 123 453.1

<151> 2001-09-28

<150> EP 02 006 712.0

<151> 2002-03-22

<150> EP 02 016 772.2

<151> 2002-07-26

<160> 217

<170> PatentIn Ver. 2.1

<210> 1

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 1

tactatacaa cctactacct caattgcc

29

<210> 2

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 2

actatgcaac ctactacctc t

21

<210> 3  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 3  
actatacaac ctctacctc a

21

<210> 4  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 4  
tggtgtttcc gcccgggaa

19

<210> 5  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 5  
tggaatgtaa agaagtatgg ag

22

<210> 6  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 6

gctcctcaaa gctggctgtg ata

23

<210> 7

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 7

tgagacacac tttgcccagt ga

22

<210> 8

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 8

tcaatggttg tctagcttta t

21

<210> 9

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 9

catatcaciaa cgatcgttcc ttt

23

<210> 10

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 10

aaaaagaaca gccactgtga ta

22

<210> 11

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 11

tggaagacta gtgattttgt tgt

23

<210> 12

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 12

gacatcttta cctgacagta tta

23

<210> 13

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 13

tcatacagct agataaccaa aga

23



<210> 14

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 14

acaaattcgg atctacaggg t

21

<210> 15

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 15

gcaagaactc agactgtgat g

21

<210> 16

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 16

accagtacct gatgtaatac tca

23

<210> 17

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

## Oligonucleotide

&lt;400&gt; 17

actcgtcaaa atggctgtga ta

22

&lt;210&gt; 18

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
Oligonucleotide

&lt;400&gt; 18

taggagagag aaaaagactg a

21

&lt;210&gt; 19

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
Oligonucleotide

&lt;400&gt; 19

tagcagcaca taatggtttg t

21

&lt;210&gt; 20

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
Oligonucleotide

&lt;400&gt; 20

gccaatattt acgtgctgct a

21

&lt;210&gt; 21

&lt;211&gt; 22

&lt;212&gt; DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 21

tacaagtgcc ttactgcag ta

22

<210> 22

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 22

tatctgcact agatgcacct ta

22

<210> 23

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 23

tcagttttgc atagatttgc aca

23

<210> 24

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 24

tacctgcact ataagcactt ta

22

<210> 25  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 25  
 tcaacatcag tctgataagc ta 22

<210> 26  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 26  
 acagttcttc aactggcagc tt 22

<210> 27  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 27  
 ggaaatccct ggcaatgtga t 21

<210> 28  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 28  
ctgttcctgc tgaactgagc ca 22

<210> 29  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 29  
tcagaccgag acaagtgcaa tg 22

<210> 30  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 30  
agcctatcct ggattacttg aa 22

<210> 31  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 31  
agcgggaactt agccactgtg aa 22

<210> 32  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 32

ctcaatagac tgtgagctcc tt

22

<210> 33

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 33

aaccgatttc agatggtgct ag

22

<210> 34

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 34

gctgcaaaca tccgactgaa ag

22

<210> 35

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 35

cagctatgcc agcatcttgc ct

22

<210> 36  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 36  
 gcaacttagt aatgtgcaat a 21

<210> 37  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 37  
 tgcaatgcaa ctacaatgca cc 22

<210> 38  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 38  
 ctccatactt ctttacattc ca 22

<210> 39  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide

<400> 39  
gctgagtgta ggatgtttac a 21

<210> 40  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 40  
gcttccagtc gaggatgttt aca 23

<210> 41  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 41  
cgcaaggctc gttctacggg tg 22

<210> 42  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 42  
tcagttatca cagtactgta 20

<210> 43  
<211> 23  
<212> DNA  
<213> Artificial Sequence



<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 43

acaaacacca ttgtcacact cca

23

<210> 44

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 44

tggcattcac cgcgtagcctt a

21

<210> 45

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 45

cacaggttaa aggtctcag gga

23

<210> 46

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 46

tcacaagtta ggtctcagg ga

22

<210> 47

<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 47  
agccaagctc agacggatcc ga

22

<210> 48  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 48  
aaaagagacc ggttcactct ga

22

<210> 49  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 49  
gcaagcccag accgaaaaa g

21

<210> 50  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 50

gcccttttaa cattgcactc

20

<210> 51

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 51

actttcgggtt atctagcttt a

21

<210> 52

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 52

acgaccatgg ctgtagactg tta

23

<210> 53

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Combined DNA/RNA Molecule:  
Oligonucleotide

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 53

tgagctacag tgcttcatct ca

22

<210> 54

<211> 18

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 54  
uuuaaccgcg aattccag 18

<210> 55  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 55  
acggaattcc tcactaaa 18

<210> 56  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 56  
cacaggtaa aggtctcag gga 23

<210> 57  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide

<400> 57  
cagccaacgg aattcctcac taaa 24

<210> 58  
<211> 22  
<212> RNA  
<213> D. melanogaster

<400> 58  
uggaauguaa agaaguaugg ag 22

<210> 59  
<211> 23  
<212> RNA  
<213> D. melanogaster

<400> 59  
uaucacagcc agcuuugaug agc 23

<210> 60  
<211> 23  
<212> RNA  
<213> D. melanogaster

<400> 60  
uaucacagcc agcuugagg agc 23

<210> 61  
<211> 22  
<212> RNA  
<213> D. melanogaster

<400> 61  
ucacugggca aagugugucu ca 22

<210> 62  
<211> 21  
<212> RNA  
<213> D. melanogaster

<400> 62  
aauaaagcuag acaaccauug a 21

<210> 63

<211> 23  
 <212> RNA  
 <213> D. melanogaster

<400> 63  
 aaaggaacga ucguugugau aug 23

<210> 64  
 <211> 22  
 <212> RNA  
 <213> D. melanogaster

<400> 64  
 uaucacagug gcuguucuuu uu 22

<210> 65  
 <211> 23  
 <212> RNA  
 <213> D. melanogaster

<400> 65  
 uggaagacua gugauuuugu ugu 23

<210> 66  
 <211> 23  
 <212> RNA  
 <213> D. melanogaster

<400> 66  
 uauuacuguc agguaaagau guc 23

<210> 67  
 <211> 23  
 <212> RNA  
 <213> D. melanogaster

<400> 67  
 ucuuugguua ucuagcugua uga 23

<210> 68  
 <211> 21  
 <212> RNA  
 <213> D. melanogaster

<400> 68  
accuguaga uccgaauug u 21

<210> 69  
<211> 21  
<212> RNA  
<213> D. melanogaster

<400> 69  
caucacaguc ugaguucuug c 21

<210> 70  
<211> 23  
<212> RNA  
<213> D. melanogaster

<400> 70  
ugaguauuac aucagguacu ggu 23

<210> 71  
<211> 22  
<212> RNA  
<213> D. melanogaster

<400> 71  
uauacagcc auuuugacga gu 22

<210> 72  
<211> 22  
<212> RNA  
<213> D. melanogaster

<400> 72  
uauacagcc auuuugauga gu 22

<210> 73  
<211> 21  
<212> RNA  
<213> D. melanogaster

<400> 73  
ucagucuuuu ucucucuccu a 21

<210> 74  
<211> 22  
<212> RNA  
<213> D. melanogaster

<400> 74  
ugagguagua gguuguauag uu 22

<210> 75  
<211> 22  
<212> RNA  
<213> Human

<400> 75  
ugagguagua gguuguauag uu 22

<210> 76  
<211> 22  
<212> RNA  
<213> Human

<400> 76  
ugagguagua gguugugugg uu 22

<210> 77  
<211> 22  
<212> RNA  
<213> Human

<400> 77  
ugagguagua gguuguaugg uu 22

<210> 78  
<211> 21  
<212> RNA  
<213> Human

<400> 78  
agagguagua gguugcauag u 21

<210> 79



<211> 21  
<212> RNA  
<213> Human

<400> 79  
ugagguagga gguuguauag u 21

<210> 80  
<211> 22  
<212> RNA  
<213> Human

<400> 80  
ugagguagua gauuguauag uu 22

<210> 81  
<211> 22  
<212> RNA  
<213> Human

<400> 81  
uagcagcaca uaaugguuug ug 22

<210> 82  
<211> 22  
<212> RNA  
<213> Human

<400> 82  
uagcagcacg uaaaauuugg cg 22

<210> 83  
<211> 20  
<212> RNA  
<213> Human

<400> 83  
acugcaguga aggcacuugu 20

<210> 84  
<211> 22  
<212> RNA  
<213> Human

<400> 84  
uaaggugcau cuagugcaga ua 22

<210> 85  
<211> 23  
<212> RNA  
<213> Human

<400> 85  
ugugcaaauc uaugcaaac uga 23

<210> 86  
<211> 23  
<212> RNA  
<213> Human

<400> 86  
ugugcaaauc caugcaaac uga 23

<210> 87  
<211> 22  
<212> RNA  
<213> Human

<400> 87  
uaaagugcuu auagugcagg ua 22

<210> 88  
<211> 22  
<212> RNA  
<213> Human

<400> 88  
uagcuauca gacugaugu ga 22

<210> 89  
<211> 22  
<212> RNA  
<213> Human

<400> 89  
aagcugccag uugaagaacu gu 22

<210> 90  
 <211> 21  
 <212> RNA  
 <213> Human

<400> 90  
 aucacauugc cagggauuuc c 21

<210> 91  
 <211> 22  
 <212> RNA  
 <213> Human

<400> 91  
 uggcucaguu cagcaggaac ag 22

<210> 92  
 <211> 22  
 <212> RNA  
 <213> Human

<400> 92  
 cauugcacuu gucucggucu ga 22

<210> 93  
 <211> 22  
 <212> RNA  
 <213> Human

<400> 93  
 uucaaguaau ccaggauagg cu 22

<210> 94  
 <211> 22  
 <212> RNA  
 <213> Human

<400> 94  
 uucaaguaau ucaggauagg uu 22

<210> 95

<211> 22  
<212> RNA  
<213> Human

<400> 95  
uucacagugg cuaaguuccg cu

22

<210> 96  
<211> 22  
<212> RNA  
<213> Human

<400> 96  
aaggagcuca cagucuauug ag

22

<210> 97  
<211> 22  
<212> RNA  
<213> Human

<400> 97  
cuagcaccau cugaaaucgg uu

22

<210> 98  
<211> 22  
<212> RNA  
<213> Human

<400> 98  
cuuucagucg gauguuugca gc

22

<210> 99  
<211> 21  
<212> RNA  
<213> Human

<400> 99  
ggcaagaugc uggcauagcu g

21

<210> 100  
<211> 21  
<212> RNA  
<213> Human

<400> 100  
uauugcacau uacuaaguug c 21

<210> 101  
<211> 19  
<212> RNA  
<213> Human

<400> 101  
gugcauugua guugcauug 19

<210> 102  
<211> 22  
<212> RNA  
<213> Human

<400> 102  
uggaauguaa agaaguauug ag 22

<210> 103  
<211> 23  
<212> RNA  
<213> Human

<400> 103  
uggaagacua gugauuuugu ugu 23

<210> 104  
<211> 23  
<212> RNA  
<213> Human

<400> 104  
ucuuugguaa ucuagcugua uga 23

<210> 105  
<211> 21  
<212> RNA  
<213> Human

<400> 105  
accuguaga uccgaauuug u 21

<210> 106  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 106  
 ugagguagua gguuguauag uu 22

<210> 107  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 107  
 ugagguagua gguugugugg uu 22

<210> 108  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 108  
 ugagguagua gguuguaugg uu 22

<210> 109  
 <211> 21  
 <212> RNA  
 <213> Mouse

<400> 109  
 agagguagua gguugcauag u 21

<210> 110  
 <211> 21  
 <212> RNA  
 <213> Mouse

<400> 110  
 ugagguagga gguuguauag u 21

<210> 111

<211> 22  
<212> RNA  
<213> Mouse

<400> 111  
ugagguagua gauuguauag uu

22

<210> 112  
<211> 22  
<212> RNA  
<213> Mouse

<400> 112  
ugagguagua guuuguacag ua

22

<210> 113  
<211> 22  
<212> RNA  
<213> Mouse

<400> 113  
ugagguagua guguguacag uu

22

<210> 114  
<211> 19  
<212> RNA  
<213> Mouse

<400> 114  
ugagguagua guuugugcu

19

<210> 115  
<211> 22  
<212> RNA  
<213> Mouse

<400> 115  
uggaauguuaa agaaguangu aa

22

<210> 116  
<211> 22  
<212> RNA  
<213> Mouse

<400> 116  
 uggaauguaa agaaguaugu ac 22

<210> 117  
 <211> 23  
 <212> RNA  
 <213> Mouse

<400> 117  
 uggaauguaa agaaguaugu auu 23

<210> 118  
 <211> 23  
 <212> RNA  
 <213> Mouse

<400> 118  
 ucuuugguua ucuagcugua uga 23

<210> 119  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 119  
 uagcagcaca uaaugguuug ug 22

<210> 120  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 120  
 uagcagcaca ucaugguuuu ca 22

<210> 121  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 121  
 uagcagcacg uaaaauuugg cg 22



<210> 122  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 122  
 uaaggugcau cuagugcaga ua 22

<210> 123  
 <211> 23  
 <212> RNA  
 <213> Mouse

<400> 123  
 ugugcaaauc caugcaaaac uga 23

<210> 124  
 <211> 23  
 <212> RNA  
 <213> Mouse

<400> 124  
 uaaagugcuu auagugcagg uag 23

<210> 125  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 125  
 uagcuuauca gacugauguu ga 22

<210> 126  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 126  
 aagcugccag uugaagaacu gu 22

<210> 127

<211> 21  
 <212> RNA  
 <213> Mouse

<400> 127  
 aucacauugc cagggauuuc c 21

<210> 128  
 <211> 23  
 <212> RNA  
 <213> Mouse

<400> 128  
 aucacauugc cagggauuac cac 23

<210> 129  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 129  
 uggcucaguu cagcaggaac ag 22

<210> 130  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 130  
 uucaaguaau ccaggauagg cu 22

<210> 131  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 131  
 uucaaguaau ucaggauagg uu 22

<210> 132  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 132  
uucacagugg cuaaguuccg cu 22

<210> 133  
<211> 20  
<212> RNA  
<213> Mouse

<400> 133  
uucacagugg cuaaguucug 20

<210> 134  
<211> 22  
<212> RNA  
<213> Mouse

<400> 134  
cuagcaccau cugaaucgg uu 22

<210> 135  
<211> 23  
<212> RNA  
<213> Mouse

<400> 135  
uagcaccauu ugaaaucagu guu 23

<210> 136  
<211> 22  
<212> RNA  
<213> Mouse

<400> 136  
uagcaccauu ugaaaucggu ua 22

<210> 137  
<211> 23  
<212> RNA  
<213> Mouse

<400> 137  
uguaaacauc cucgacugga agc 23

<210> 138  
<211> 22  
<212> RNA  
<213> Mouse

<400> 138  
cuuucagucg gauguuugca gc 22

<210> 139  
<211> 21  
<212> RNA  
<213> Mouse

<400> 139  
uguaaacauc cuacacucag c 21

<210> 140  
<211> 23  
<212> RNA  
<213> Mouse

<400> 140  
uguaaacauc cuacacucuc agc 23

<210> 141  
<211> 22  
<212> RNA  
<213> Mouse

<400> 141  
uguaaacauc cccgacugga ag 22

<210> 142  
<211> 20  
<212> RNA  
<213> Mouse

<400> 142  
accguagau ccgaucuugu 20

<210> 143

<211> 22  
<212> RNA  
<213> Mouse

<400> 143  
cacccguaga accgaccuug cg 22

<210> 144  
<211> 20  
<212> RNA  
<213> Mouse

<400> 144  
uacaguacug ugauaacuga 20

<210> 145  
<211> 23  
<212> RNA  
<213> Mouse

<400> 145  
uggaguguga caaugguguu ugu 23

<210> 146  
<211> 23  
<212> RNA  
<213> Mouse

<400> 146  
uggaguguga caaugguguu uga 23

<210> 147  
<211> 22  
<212> RNA  
<213> Mouse

<400> 147  
uggaguguga caaugguguu ug 22

<210> 148  
<211> 21  
<212> RNA  
<213> Mouse

<400> 148  
cauuauuacu uuugguacgc g 21

<210> 149  
<211> 22  
<212> RNA  
<213> Mouse

<400> 149  
uuaaggcacg cggugaaugc ca 22

<210> 150  
<211> 21  
<212> RNA  
<213> Mouse

<400> 150  
uuaaggcacg cgggugaug c 21

<210> 151  
<211> 23  
<212> RNA  
<213> Mouse

<400> 151  
ucccugagac ccuuuaaccu gug 23

<210> 152  
<211> 22  
<212> RNA  
<213> Mouse

<400> 152  
ucccugagac ccuaacuugu ga 22

<210> 153  
<211> 21  
<212> RNA  
<213> Mouse

<400> 153  
ucguaccgug aguaauaauug c 21

<210> 154  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 154  
 ucggauccgu cugagcuugg cu 22

<210> 155  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 155  
 ucacagugaa ccggucucuu uu 22

<210> 156  
 <211> 21  
 <212> RNA  
 <213> Mouse

<400> 156  
 cuuuuuucgg ucugggcuug c 21

<210> 157  
 <211> 20  
 <212> RNA  
 <213> Mouse

<400> 157  
 cagugcaaug uaaaaagggc 20

<210> 158  
 <211> 21  
 <212> RNA  
 <213> Mouse

<400> 158  
 uaaagcuaga uaaccgaaag u 21

<210> 159

<211> 23  
 <212> RNA  
 <213> Mouse

<400> 159  
 uaacagucua cagccauggu cgu 23

<210> 160  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 160  
 uugguccccu ucaaccagcu gu 22

<210> 161  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 161  
 ugugacuggu ugaccagagg ga 22

<210> 162  
 <211> 24  
 <212> RNA  
 <213> Mouse

<400> 162  
 uauggcuuuu uauuccuaug ugaa 24

<210> 163  
 <211> 23  
 <212> RNA  
 <213> Mouse

<400> 163  
 acuccauuug uuuugaugau gga 23

<210> 164  
 <211> 22  
 <212> RNA  
 <213> Mouse



<400> 164	
uauugcuuaa gaauacgcu ag	22
<210> 165	
<211> 17	
<212> RNA	
<213> Mouse	
<400> 165	
agcugguguu gugaauc	17
<210> 166	
<211> 18	
<212> RNA	
<213> Mouse	
<400> 166	
ucuacagugc acgugucu	18
<210> 167	
<211> 21	
<212> RNA	
<213> Mouse	
<400> 167	
agugguuuaa ccuauggua g	21
<210> 168	
<211> 21	
<212> RNA	
<213> Mouse	
<400> 168	
aacacugucu gguaaagaug g	21
<210> 169	
<211> 20	
<212> RNA	
<213> Mouse	
<400> 169	
cauaaaguag aaagcacuac	20

<210> 170  
<211> 22  
<212> RNA  
<213> Mouse

<400> 170  
uguaguguuu ccuacuuuau gg 22

<210> 171  
<211> 22  
<212> RNA  
<213> Mouse

<400> 171  
ugagaugaag cacuguagcu ca 22

<210> 172  
<211> 22  
<212> RNA  
<213> Mouse

<400> 172  
uacaguauag augauguacu ag 22

<210> 173  
<211> 24  
<212> RNA  
<213> Mouse

<400> 173  
guccaguuuu cccaggauc ccuu 24

<210> 174  
<211> 23  
<212> RNA  
<213> Mouse

<400> 174  
ugagaacuga auccauggg uuu 23

<210> 175

<211> 21  
 <212> RNA  
 <213> Mouse

<400> 175  
 guguguggaa augcuucugc c 21

<210> 176  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 176  
 ucagugcacu acagaacuuu gu 22

<210> 177  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 177  
 ucuggcuccg ugucuucacu cc 22

<210> 178  
 <211> 23  
 <212> RNA  
 <213> Mouse

<400> 178  
 ucucccaacc cuuguaccag ugu 23

<210> 179  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 179  
 cuagacugag gcuccuugag gu 22

<210> 180  
 <211> 21  
 <212> RNA  
 <213> Mouse

<400> 180  
 ucagugcaug acagaacuug g 21

<210> 181  
 <211> 20  
 <212> RNA  
 <213> Mouse

<400> 181  
 uugcauaguc acaaaaguga 20

<210> 182  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 182  
 uagguuaucc guguugccuu cg 22

<210> 183  
 <211> 22  
 <212> RNA  
 <213> Mouse

<400> 183  
 uuaaugcuaa uugugauagg gg 22

<210> 184  
 <211> 23  
 <212> RNA  
 <213> Human/Mouse

<400> 184  
 aacaucaac gcugucggug agu 23

<210> 185  
 <211> 22  
 <212> RNA  
 <213> Human/Mouse

<400> 185  
 uuuggcaaug guagaacuca ca 22

<210> 186

<211> 23

<212> RNA

<213> Human/Mouse

<400> 186

uauggcacug guagaauca cug

23

<210> 187

<211> 22

<212> RNA

<213> Human/Mouse

<400> 187

cuuuuugcgg ucugggcuug uu

22

<210> 188

<211> 22

<212> RNA

<213> Human/Mouse

<400> 188

uggacggaga acugauaagg gu

22

<210> 189

<211> 18

<212> RNA

<213> Human/Mouse

<400> 189

uggagagaaa ggcaguuc

18

<210> 190

<211> 23

<212> RNA

<213> Human/Mouse

<400> 190

caaagaauuc uccuuuuggg cuu

23

<210> 191

<211> 22

<212> RNA

<213> Human/Mouse

<400> 191

ucgugucuug uguugcagcc gg

22

<210> 192

<211> 21

<212> RNA

<213> Human/Mouse

<400> 192

uaacacuguc ugguaacgau g

21

<210> 193

<211> 22

<212> RNA

<213> Human/Mouse

<400> 193

caucccuugc augguggagg gu

22

<210> 194

<211> 23

<212> RNA

<213> Human/Mouse

<400> 194

gugccuacug agcugacauc agu

23

<210> 195

<211> 22

<212> RNA

<213> Human/Mouse

<400> 195

ugauauguuu gauauuuag gu

22

<210> 196

<211> 22

<212> RNA

<213> Human/Mouse

<400> 196  
caacggauc caaaagcag cu 22

<210> 197  
<211> 18  
<212> RNA  
<213> Human/Mouse

<400> 197  
cugaccuaug aaugaca 18

<210> 198  
<211> 22  
<212> RNA  
<213> Human/Mouse

<400> 198  
uaccacaggg uagaaccacg ga 22

<210> 199  
<211> 21  
<212> RNA  
<213> Human/Mouse

<400> 199  
aacuggccua caaagucca g 21

<210> 200  
<211> 22  
<212> RNA  
<213> Human/Mouse

<400> 200  
uguaacagca acuccaugug ga 22

<210> 201  
<211> 21  
<212> RNA  
<213> Human/Mouse

<400> 201  
uagcagcaca gaaauauugg c 21

<210> 202

<211> 21

<212> RNA

<213> Human/Mouse

<400> 202

uagguaguuu cauguuguug g

21

<210> 203

<211> 22

<212> RNA

<213> Human/Mouse

<400> 203

uucaccaccu ucuccaccca gc

22

<210> 204

<211> 19

<212> RNA

<213> Human/Mouse

<400> 204

gguccagagg ggagauagg

19

<210> 205

<211> 22

<212> RNA

<213> Human/Mouse

<400> 205

cccaguguuc agacuaccug uu

22

<210> 206

<211> 23

<212> RNA

<213> Human/Mouse

<400> 206

uaauacugcc ugguaaugau gac

23

<210> 207



<211> 21

<212> RNA

<213> Human/Mouse

<400> 207

uacucaguuaa ggcauuguuc u

21

<210> 208

<211> 22

<212> RNA

<213> Human/Mouse

<400> 208

agagguaauag cgcaugggaa ga

22

<210> 209

<211> 21

<212> RNA

<213> Human/Mouse

<400> 209

ugaaauguuu aggaccacua g

21

<210> 210

<211> 23

<212> RNA

<213> Human/Mouse

<400> 210

uucccuuugu cauccuaugc cug

23

<210> 211

<211> 22

<212> RNA

<213> Human/Mouse

<400> 211

uccuucuuuc caccggaguc ug

22

<210> 212

<211> 23

<212> RNA

<213> Human/Mouse

<400> 212  
gugaaauguu uaggaccacu aga 23

<210> 213  
<211> 22  
<212> RNA  
<213> Human/Mouse

<400> 213  
uggaauguaa ggaagugugu gg 22

<210> 214  
<211> 22  
<212> RNA  
<213> Human/Mouse

<400> 214  
uacaguaguc ugcacauugg uu 22

<210> 215  
<211> 22  
<212> RNA  
<213> Human/Mouse

<400> 215  
cccuguagaa ccgaauugu gu 22

<210> 216  
<211> 24  
<212> RNA  
<213> Human/Mouse

<400> 216  
aaccguaga uccgaacuug uгаа 24

<210> 217  
<211> 23  
<212> RNA  
<213> Human/Mouse

<400> 217  
gcuuccucug gcucuccuc cuc 23

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- BLACK BORDERS
- IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT OR DRAWING
- BLURRED OR ILLEGIBLE TEXT OR DRAWING
- SKEWED/SLANTED IMAGES
- COLOR OR BLACK AND WHITE PHOTOGRAPHS
- GRAY SCALE DOCUMENTS
- LINES OR MARKS ON ORIGINAL DOCUMENT
- REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**